

**FT-IV-KVPY-CLASS-XII**  
**FULL TEST – IV**

**PART – I**  
**MATHEMATICS**

1. Let  $a_n = \sum_{k=1}^n \frac{1}{k(n+1-k)}$ , then for  $n \geq 2$
- (A)  $a_{n+1} > a_n$  (B)  $a_{n+1} < a_n$   
(C)  $a_{n+1} = a_n$  (D)  $a_{n+1} - a_n = \frac{1}{n}$

Ans. B

Sol. We have  $a_n = \sum_{k=1}^n \left( \frac{1}{k} + \frac{1}{n+1-k} \right)$

$$= \frac{2}{n+1} \sum_{k=1}^n \left( \frac{1}{k} + \frac{1}{n+1-k} \right)$$
$$= \frac{2}{n+1} \sum_{k=1}^n \frac{1}{k}$$

For  $n \geq 2$

$$\frac{1}{2}(a_n - a_{n+1}) = \frac{1}{n+1} \sum_{k=1}^n \frac{1}{k} - \frac{1}{n+2} \sum_{k=1}^{n+1} \frac{1}{k}$$
$$= \left( \frac{1}{n+1} - \frac{1}{n+2} \right) \sum_{k=1}^n \frac{1}{k} - \frac{1}{(n+1)(n+2)}$$
$$= \frac{1}{(n+1)(n+2)} \sum_{k=2}^n \frac{1}{k} > 0$$

$\Rightarrow a_n > a_{n+1}$

2. If  $\cos\theta_1 + 2 \cos\theta_2 + 3 \cos\theta_3 = 6$  then  $\tan\theta_1 + \tan\theta_2 + \tan\theta_3$  equals to
- (A) 1/2 (B) 6  
(C) 0 (D) 3

Ans. C

Sol. Since,  $\cos\theta \leq 1$ .

If  $\cos\theta_1 + 2\cos\theta_2 + \sec\theta_3 = 6$ , then

$$\cos\theta_1 = \cos\theta_2 = \cos\theta_3 = 1 \Rightarrow \tan\theta_1 = \tan\theta_2 = \tan\theta_3 = 0$$

3. The number of non – negative integral solutions of  $x_1 + x_2 + x_3 + 4x_4 = 20$  is
- (A) 530 (B) 532  
(C) 534 (D) 536

Ans. D

Sol. Coefficient of  $x^{20}$  in  $(1-x)^{-3}(1-x^4)^{-1}$

Coefficient of  $x^2$  in

$$\begin{aligned} & (1 + {}^3C_1x + {}^4C_2x^2 + \dots)(1 + x^4 + x^8 + x^{12} + x^{16} + x^{20}) \\ &= 1 + {}^6C_4 + {}^{10}C_8 + {}^{14}C_{12} + {}^{18}C_{16} + {}^{22}C_{20} \\ &= 1 + {}^6C_2 + {}^{10}C_2 + {}^{14}C_2 + {}^{18}C_2 + {}^{21}C_2 \\ &= 1 + 15 + 45 + 91 + 153 + 231 \\ &= 536 \end{aligned}$$

4. Let AB be a line segment of length 4 unit with the point A on the line  $y = 2x$  and B on the line  $y = x$ . Then locus of middle point of all such line segment is

- (A) a parabola (B) an ellipse  
(C) a hyperbola (D) a circle

Ans. B

Sol. Let  $B = (\alpha, \alpha)$  and middle point AB is  $(h, k)$

Then,  $A = (2h - \alpha, 2k - \alpha)$

lies on  $y = 2x$

then,  $(2k - \alpha) = 2(2h - \alpha) \dots(i)$

$\therefore \alpha = 4h - 2k$

$\Rightarrow \frac{|AB| = 4}{\sqrt{(2h - 2\alpha)^2 + (2k - 2\alpha)^2}} = 4$

or  $(h - \alpha)^2 + (k - \alpha)^2 = 4$

or  $[h - (4h - 2k)]^2 + [k - (4h - 2k)]^2 = 4$


$\Rightarrow (-3h + 2k)^2 + (-4h + 3k)^2 = 4$

or  $25h^2 + 13k^2 - 36hk = 4$

Required locus is  $25x^2 + 13y^2 - 36xy - 4 = 0$

Here,  $h^2 < ab$  and  $\Delta \neq 0$

$\therefore$  ellipse



5. If  $(1+x)^n = \sum_{r=0}^n a_r x^r$  and  $b_r = 1 + \frac{a_r}{a_{r-1}}$  and  $\prod_{r=1}^n b_r = \frac{(101)^{100}}{100!}$ , then n is

- (A) 99 (B) 100  
(C) 101 (D) 102

Ans. B

Sol.  $(1+x)^n = \sum_{r=0}^n {}^nC_r x^r = \sum_{r=0}^n a_r x^r$  (given)

$$\begin{aligned} \therefore a_r &= {}^n C_r \\ \text{Also, } b_r &= 1 + \frac{a_r}{a_{r-1}} = \frac{a_r}{a_{r-1}} = 1 + \frac{{}^n C_r}{{}^n C_{r-1}} = \frac{{}^{n+1} C_r}{{}^n C_{r-1}} \\ b_r &= \binom{n+1}{r} \\ \therefore \prod_{r=1}^n b_r &= \prod_{r=2}^n \binom{n+1}{r} = \frac{(n+1)^n}{n!} \\ &= \frac{(101)^{100}}{100!} \quad (\text{given}) \\ \therefore n &= 100 \end{aligned}$$

6. The line which intersect the skew lines  $y = mx, z = c$ ;  $y = -mx, z = -c$  and the  $x$  - axis lie on the surface
- (A)  $cz = mxy$  (B)  $cy = mxz$   
 (C)  $xy = cmz$  (D) none of these

Ans. B

Sol. Equation of the planes through  $y = mx, z = c$  and  $y = -mx, z = -c$  are respectively

$$(y - mx) + \lambda_1(z - c) = 0 \quad \dots\dots(i)$$

$$\text{and } (y + mx) + \lambda_2(z + c) = 0 \quad \dots\dots(ii)$$

It meets at  $x$  - axis i.e.  $y = 0 = z$

$$\therefore \lambda_2 = \lambda_1$$

From equation (i) and (ii),

$$\frac{y - mx}{z - c} = \frac{y + mx}{z + c}$$

$$\therefore cy = mzx$$

7. The number of six digit numbers that can be formed from the digits 1, 2, 3, 4, 5, 6 and 7, so that digits do not repeat and the terminal digits are even is
- (A) 144 (B) 72  
 (C) 288 (D) 720

Ans. D

Sol. Terminal digits are the first and last digits.

$\therefore$  Terminal digits are even

$\therefore$  1<sup>st</sup> place can be filled in 3 ways and last place can be filled in 2<sup>nd</sup> ways and remaining places can be filled in  ${}^5 P_4 = 120$  ways

Hence, the number of six digit numbers, the terminal digits are even, is  
 $= 3 \times 120 \times 2 = 720$

8. Three players A, B, C in this order, cut a pack of cards, and the whole pack is reshuffled after each cut. If the winner is one who first draws a diamond, then C's chance of winning is

- (A)  $\frac{9}{28}$  (B)  $\frac{9}{37}$   
 (C)  $\frac{9}{64}$  (D)  $\frac{27}{64}$

Ans. B

Sol. Probability of diamond card =  $\frac{{}^{13}C_1}{{}^{52}C_1} = \frac{1}{4}$

Probability of C's winning

$$= P(\bar{A})P(\bar{B})P(C) + P(\bar{A})P(\bar{B})P(\bar{C})P(\bar{A})P(\bar{B})P(C) + \dots$$

$$= \frac{P(\bar{A})P(\bar{B})P(\bar{C})}{1 - P(\bar{A})P(\bar{B})P(\bar{C})} = \frac{\frac{3}{4} \times \frac{3}{4} \times \frac{1}{4}}{1 - \frac{3}{4} \times \frac{3}{4} \times \frac{3}{4}} = \frac{9}{37}$$

9. A three digit number, which is multiple of 11, is chosen at random. The probability that the number so chosen is also a multiple of 9 is equal to

- (A)  $\frac{1}{9}$  (B)  $\frac{2}{9}$   
 (C)  $\frac{1}{100}$  (D)  $\frac{9}{100}$

Ans. A

Sol. The number of three digit numbers, which are multiple 11 =  $90 - 9 = 81$ . Again the number, which are divisible of 9 also, are divisible by 99, whose number is  $10 - 1 = 9$ .

So, required probability =  $\frac{9}{81} = \frac{1}{9}$ .

10. If x, y, z are integers in AP, lying between 1 and 9 and x51, y41 and z31 are three digit

numbers, then the value of  $\begin{vmatrix} 5 & 4 & 3 \\ x51 & y41 & z31 \\ x & y & z \end{vmatrix}$  is

- (A)  $x + y + z$  (B)  $x - y + z$   
 (C) 0 (D)  $x + 2y + z$

Ans. C

Sol. Let  $\Delta = \begin{vmatrix} 5 & 4 & 3 \\ x51 & y41 & z31 \\ x & y & z \end{vmatrix}$

$$\Delta = \begin{vmatrix} 5 & 4 & 3 \\ 100x + 50 + 1 & 100y + 40 + 1 & 100z + 30 + 1 \\ x & y & z \end{vmatrix}$$

Applying  $R_2 \rightarrow R_2 - (100R_3 + 10R_1)$ , then

$$\Delta = \begin{vmatrix} 5 & 4 & 3 \\ 1 & 1 & 1 \\ x & y & z \end{vmatrix}$$

$$\text{Applying } C_2 \rightarrow C_2 - \frac{1}{2}(C_1 + C_3), \text{ then } \Delta = \begin{vmatrix} 5 & 0 & 3 \\ 1 & 0 & 1 \\ x & y - \frac{1}{2}(x+z) & z \end{vmatrix}$$

$$= \begin{vmatrix} 5 & 0 & 3 \\ 1 & 0 & 1 \\ x & 0 & z \end{vmatrix} = 0 \quad (\because x, y, z \text{ are in AP})$$

11. For the equations;  $x + 2y + 3z = 1$ ,  $2x + y + 3z = 2$ ,  $5x + 5y + 9z = 4$   
 (A) there is only one solution                      (B) there exists infinitely many solutions  
 (C) there is no solution                              (D) none of the above

Ans. A

Sol.  $\therefore \begin{vmatrix} 1 & 2 & 3 \\ 2 & 1 & 3 \\ 5 & 5 & 9 \end{vmatrix} = 1(9 - 15) - 2(18 - 15) + 3(10 - 5)$   
 $= -6 - 6 + 15 = 3 \neq 0$   
 i.e. only one solution.

12.  $f(x) = 1 + \sin x [\cos x]$ ,  $0 < x \leq \frac{\pi}{2}$  ([.] denotes the greatest integer function)  
 (A) is continuous in  $\left(0, \frac{\pi}{2}\right)$                       (B) is strictly decreasing in  $\left(0, \frac{\pi}{2}\right)$   
 (C) is strictly increasing in  $\left(0, \frac{\pi}{2}\right)$                       (D) has global maximum value 2

Ans. A

Sol.  $\therefore 0 < x < \frac{\pi}{2}$                        $\therefore 0 \leq \cos x < 1$  then  $[\cos x] = 0$   
 $\therefore f(x) = 1$   
 Hence,  $f(x)$  is continuous in  $\left(0, \frac{\pi}{2}\right)$

13. The solution of the differential equation  $\frac{dy}{dx} = \frac{1}{xy(x^2 \sin y^2 + 1)}$  is
- (A)  $x^2(\cos y^2 - \sin y^2 - ce^{-y^2}) = 2$                       (B)  $y^2(\cos x^2 - \sin y^2 - ce^{-y^2}) = 2$   
 (C)  $x^2(\cos y^2 - \sin y^2 - e^{-y^2}) = 4c$                       (D) none of these

Ans. A

Sol.  $\frac{dy}{dx} = \frac{1}{xy(x^2 \sin y^2 + 1)} \Rightarrow \frac{dx}{dy} = xy(x^2 \sin y^2 + 1)$   
 $\Rightarrow \frac{1}{x^3} \frac{dx}{dy} - \frac{1}{x^2} y = y \sin y^2 \dots\dots\dots(1)$

Put  $-\frac{1}{x^2} = u$ .  
 $\Rightarrow \frac{du}{dy} + 2uy = 2y \sin y^2$

This is a linear differential equation.

14.  $\int \frac{\operatorname{cosec}^2 x - 2005}{\cos^{2005} x} dx$  is equal to
- (A)  $\frac{\cot x}{(\cos x)^{2005}} + C$                       (B)  $\frac{\tan x}{(\cos x)^{2005}} + C$   
 (C)  $-\frac{\tan x}{(\cos x)^{2005}} + C$                       (D)  $-\frac{\cot x}{(\cos x)^{2005}} + C$

Ans. D

Sol.  $I = \int \frac{\operatorname{cosec}^2 x - 2005}{\cos^{2005} x} dx$   
 $= \int (\cos x)^{-2005} \operatorname{cosec}^2 x dx - 2005 \int \frac{dx}{\cos^{2005} x}$   
 $= (\cos x)^{-2005} (-\cot x) - \int (-2005)(\cos x)^{-2006} (-\sin x)(-\cot x) dx - 2005 \int \frac{dx}{\cos^{2005} x}$   
 $= -\frac{\cot x}{(\cos x)^{2005}} + C$

15. Number of common tangents with finite slope to the curves  $xy = c^2$  and  $y^2 = 4ax$  is
- (A) 0    (B) 1  
 (C) 2    (D) 4

Ans. B

Sol. Equation of PQ is  $yt - x = at^2$  .....(1)

$$\frac{x}{t'} + yt' = 2c \quad \text{.....(2)}$$


$\therefore$  (1) and (2) are identical

$$\Rightarrow -t' = \frac{t}{t'} = \frac{at^2}{2c}$$

$$\Rightarrow \frac{a(t')^4}{2c} = -t'$$

$$\Rightarrow (t')^3 = -\frac{2c}{a}$$

$\Rightarrow$  Only one real value of  $t'$  exists.



16. If the roots of  $x^5 - 40x^4 + Px^3 + Qx^2 + Rx + S = 0$  are in G.P. and the sum of their reciprocals is 10, then  $|S|$  is equal to

- (A) 4  
(C) 8

- (B) 6  
(D) 32

Ans. D

Sol. The roots of the equation  $x^5 - 40x^4 + Px^3 + Qx^2 + Rx + S = 0$  are in G.P. Let the roots be  $a, ar, ar^2, ar^3, ar^4$ .

$$\therefore a + ar + ar^2 + ar^3 + ar^4 = 40$$

$$\text{and } \frac{1}{a} + \frac{1}{ar} + \frac{1}{ar^2} + \frac{1}{ar^3} + \frac{1}{ar^4} = \frac{1+r+r^2+r^3+r^4}{ar^4} = 10$$

$$\Rightarrow a^2r^4 = 4 \text{ or } ar^2 = \pm 2$$

$$\text{Now, } -S = \text{product of roots} = a^5r^{10} = (ar^2)^5 = \pm 32$$

$$\therefore |S| = 32.$$

17. Let  $f(x) = -x^3 + px^2 + qx + 6 \operatorname{sgn}(x^2 + x + 1)$ , where  $p, q \in \mathbb{R}$  and  $\operatorname{sgn}$  stands for signum function. If the largest possible interval in which  $f'(x)$  is positive is  $\left(-\frac{5}{3}, 1\right)$ , then  $(p+q)$

equals

- (A) 6  
(C) -4

- (B) 4  
(D) -6

Ans. B

Sol.  $f(x) = -x^3 + px^2 + qx + 6$

$$f'(x) = -3x^2 + 2px + q > 0$$

$$3x^2 - 2px - q < 0$$

$$x^2 - \frac{2p}{3}x - \frac{q}{3} < 0$$

Given,  $\left(x - \left(-\frac{5}{3}\right)\right)(x - 1) < 0$

$$x^2 + \frac{2}{3}x - \frac{5}{3} < 0$$

$$p = -1; q = 5$$

18. If the line  $x + y - 1 = 0$  is a tangent to a parabola with focus (1, 2) at A and intersects the directrix at B and the tangent at the vertex at C, then AC. BC is equal to

(A) 2

(B) 1

(C)  $\frac{1}{2}$

(D)  $\frac{1}{4}$

Ans. A

Sol.  $(BC)(AC) = (CS)^2$

19. If  $\alpha, \beta, \gamma$  are the angle of a triangle and the system of equations

$$\cos(\alpha - \beta)x + \cos(\beta - \gamma)y + \cos(\gamma - \alpha)z = 0$$

$$\cos(\alpha + \beta)x + \cos(\beta + \gamma)y + \cos(\gamma + \alpha)z = 0$$

$$\sin(\alpha + \beta)x + \sin(\beta + \gamma)y + \sin(\gamma + \alpha)z = 0$$

has non-trivial solutions, then the triangle is necessarily

(A) equilateral

(B) isosceles

(C) right angled

(D) acute angled

Ans. B

Sol. Let  $\Delta = \begin{vmatrix} \cos(\alpha - \beta) & \cos(\beta - \gamma) & \cos(\gamma - \alpha) \\ \cos(\alpha + \beta) & \cos(\beta + \gamma) & \cos(\gamma + \alpha) \\ \sin(\alpha + \beta) & \sin(\beta + \gamma) & \sin(\gamma + \alpha) \end{vmatrix}$

It is clear that either  $\alpha = \beta$  or  $\beta = \gamma$  or  $\gamma = \alpha$  is sufficient to make  $\Delta = 0$ . It is not necessary that the triangle be equilateral.

20. Let  $f : [0, 1] \rightarrow \mathbb{R}$  be a continuous function and assumes only rational values. If  $f(0) = 2$ ,

then the value of  $\tan^{-1}\left(f\left(\frac{1}{2}\right)\right) + \tan^{-1}\left(\frac{3}{2}f\left(\frac{1}{2}\right)\right)$  is

(A)  $\frac{\pi}{4}$

(B)  $\frac{\pi}{6}$

(C)  $\frac{3\pi}{4}$

(D)  $\pi$

Ans. C



Sol.  $f(0) = 2 \Rightarrow f\left(\frac{1}{2}\right) = 2$

Required expression =  $\tan^{-1}(2) + \tan^{-1}\left(\frac{3}{2} \cdot 2\right)$

=  $\tan^{-1}(2) + \tan^{-1}(3) = \pi + \tan^{-1}\left(\frac{2+3}{1-2 \cdot 3}\right)$

=  $\pi - \tan^{-1}(1) = \pi - \frac{\pi}{4} = \frac{3\pi}{4}$

## PHYSICS


21. A block of mass  $m$  is placed on the top of a wedge having undefined smooth curved surface. If the wedge is now accelerated horizontally with acceleration  $a$ , then the speed of block with respect to wedge when it reaches the bottom of wedge is

(A)  $\sqrt{2gh}$

(B)  $\sqrt{2(a+g)h}$

(C)  $\sqrt{2(al-gh)}$

(D)  $\sqrt{2(al+gh)}$



Ans. D

Sol.  $ma\ell + mgh = \frac{1}{2}mv^2$

$\therefore v = \sqrt{2(al+gh)}$


22. At an instant  $m_1$  and  $m_2$  are having speed  $v_1$  and  $v_2$  and acceleration  $a_1$  and  $a_2$ . Then

(A)  $v_1 = v_2 \sec \theta$

(B)  $a_1 = a_2 \cos \theta$  only at an instant when the blocks starts from rest

(C)  $a_1 = a_2 \cos \theta$  for all instant

(D)  $v_1 = v_2 \sin \theta$



Ans. B

Sol. differentiating the relation  $v_1 = v_2 \cos \theta$  with respect to time.

23. A particle is projected with speed  $u$  at an angle  $\theta$  with ground. The radius of curvature at highest point is

(A)  $\frac{u^2 \cos^2 \theta}{g}$

(B)  $\frac{u^2 \sin^2 \theta}{g}$

(C)  $\frac{u^2 \cos^2 \theta}{2g}$

(D)  $\frac{u^2 \sin^2 \theta}{2g}$

Ans. A

Sol.  $R = \frac{v_{\text{net}}^2}{a_r} \Rightarrow \vec{a}_r \perp \vec{v}_{\text{net}}$

24. If the change in the value of 'g' at a height h above the surface of the earth is the same as a depth x below it, then (both x and h being much smaller than the radius of the earth):
- (A)  $x = h$  (B)  $x = 2h$   
 (C)  $x = 1/2 h$  (D)  $x = h^2$ .

Ans. B

Sol. Above surface of earth at height  $h \ll R$

$$g' = g \left[ 1 - \frac{2h}{R} \right] \Rightarrow \frac{\Delta g}{g} = \frac{2h}{R}$$

surface below earth  $x \ll R$ .



$$g' = g \left[ 1 - \frac{x}{R} \right] \Rightarrow \frac{\Delta g}{g} = \frac{x}{R}$$

25. A system consists of two stars of equal masses that revolve in a circular orbit about a centre of mass midway between them. Orbital speed of each star is v & period is T. Find the mass M of each star: (G is gravitational constant)
- (A)  $\frac{2Gv^3}{\pi T}$  (B)  $\frac{v^3 T}{\pi G}$   
 (C)  $\frac{v^3 T}{2\pi G}$  (D)  $\frac{2Tv^3}{\pi G}$

Ans. D

Sol.  $\frac{GM^2}{(2r)^2} = \frac{MV^2}{r}$  Or  $M = \frac{4v^2 r}{G}$  &  $T = \frac{2\pi r}{v}$ ,  $r = \frac{Tv}{2\pi}$

26. A sphere of mass M and radius b has a concentric cavity of radius a as shown in figure. The graph showing variation of gravitational potential V with distance r from the center of sphere is




Ans. B

Sol. Gravitational potential inside a shell = constant

27. In the figure shown, the pulley is light, wedge is fixed and all the surfaces are smooth. Tension in the string is

- (A)  $mg \sin \theta$  (B)  $\frac{mg \sin \theta}{2}$   
 (C)  $mg \frac{\sin \theta}{3}$  (D)  $mg$




Ans. B

Sol.  $mg \sin \theta - T = ma$   
 $T = ma$

28. Air is blowing and is providing a constant horizontal acceleration  $a_x = g$  to the particle as shown in the figure. Particle is projected from point P with a velocity  $u$  in upward direction. Let Q be the highest point of particle. Speed of the particle at highest point Q is

- (A)  $\sqrt{2}u$  (B)  $u$   
 (C)  $u/\sqrt{2}$  (D) None



Ans. B

Sol.  $v_x = u_x + a_x t$   
 $v_x = 0 + gt$  ..... (1)  
 $v_y = u_y + a_y t$   
 $0 = u - gt$   
 $t = \frac{u}{g}$  .....(2)

By (1) and (2) we get

$$v_x = u \text{ and } v_y = 0$$

Hence net velocity =  $u$

29. When the current in a coil changes from 8A to 2A in  $3 \times 10^{-2}$  seconds, the e.m.f induced in the coil is 2 volts. The self inductance of the cell in millihenry is

- (A) 1 (B) 5  
 (C) 20 (D) 10

Ans. D

Sol.  $\varepsilon = L \frac{dl}{dt} \Rightarrow 2 = L \times \frac{6A}{3 \times 10^{-2}}$   
 $L = 10^{-2}$  Henry = 10 mH

30. Electric charge  $q$ ,  $q$  and  $-2q$  are placed at the corners of an equilateral triangle ABC of side  $L$ . The magnitude of electric dipole moment of the system is:  
 (A)  $qL$  (B)  $2qL$   
 (C)  $\sqrt{3}qL$  (D)  $4qL$

Ans. C

Sol. Two dipoles will be formed at angle  $60^\circ$  to each other.

$$\therefore P_{\text{net}} = \sqrt{(qL)^2 + (qL)^2 + 2(qL)^2 \cos 60^\circ} = \sqrt{3}qL$$

31. A uniform magnetic field of  $30 \text{ mT}$  exists in the  $+X$  direction. A particle of charge  $+e$  and mass  $1.67 \times 10^{-27} \text{ kg}$  is projected through the field in the  $+Y$  direction with a speed of  $4.8 \times 10^6 \text{ m/s}$ . Radius of the circular path followed by the particle is  
 (A)  $6.17 \text{ m}$  (B)  $1.67 \text{ m}$   
 (C)  $1.76 \text{ m}$  (D)  $1.77 \text{ m}$

Ans. B

Sol. (B)  $F = qVB \sin\theta$   
 $= (1.6 \times 10^{-19}) (4.8 \times 10^6) (30 \times 10^{-3}) \sin 90^\circ$   
 $= 230.4 \times 10^{-6} \text{ N}$ .

The direction of the force is in the  $(-z)$  direction.


(b) If the particle were negatively charged, the magnitude of the force will be the same but the direction will be along  $(+z)$  direction.

(c) As  $V \perp B$ , the path described is a circle

$$R = mV/qB$$

$$= (1.67 \times 10^{-27}) \cdot (4.8 \times 10^6) / (1.6 \times 10^{-19}) \cdot (30 \times 10^{-3}) = 1.67 \text{ m}.$$

32. The effective resistance between the point A and B will be  
 (A)  $4 \Omega$   
 (B)  $2 \Omega$   
 (C)  $6 \Omega$   
 (D)  $8 \Omega$



Ans. B

Sol. Resistors AF and FE are in series with each other. Therefore, network AEF reduces to a parallel combination of two resistors of  $6 \Omega$  each.

$$R_{\text{eq}} = \frac{6 \times 6}{6 + 6} = 3 \Omega.$$

Similarly, the resistance between A and D is given

$$\frac{6 \times 6}{6 + 6} = 3 \Omega.$$

Now, resistor AC is in parallel with the series combination of AD and DC. Therefore, the resistance between A and C is  $\frac{6 \times 6}{6 + 6} = 3\Omega$ .


AC + CB = 3 + 3 = 6  $\Omega$ . Since they are in series resistance between A and B is given by  $\frac{1}{R} = \frac{1}{6} + \frac{1}{3} = \frac{3}{6}$  or  $R_{AB} = 2\Omega$ .

33. Potential in the x-y plane is given as  $V = 5(x^2 + xy)$  volts. The electric field at the point (1, -2) will be
- (A)  $3\hat{j}$  V/m (B)  $-5\hat{j}$  V/m  
 (C)  $5\hat{j}$  V/m (D)  $-3\hat{j}$  V/m

Ans. B

Sol.  $E_x = -\frac{\partial V}{\partial x} = -(10x + 5y) = -10 + 10 = 0$   
 $E_y = -\frac{\partial V}{\partial y} = -5x = -5$   
 $\therefore \vec{E} = -5\hat{j}$  V/m.


34. The equivalent capacitance between points A and B of the circuit will be
- (A) 12C (B) 6C  
 (C) 3C (D) 24C




Ans. C

Sol. Equivalent circuit of the above figure can be drawn as

$C_{AB} = 3C$



35. The wire loop PQRSP formed by joining two semi-circular wires of radii  $R_1$  and  $R_2$  carries a current  $I$  as shown in figure. What is the magnetic induction at the centre O and magnetic moment of the loop?



- (A)  $\frac{\pi I}{2}(R_2^2 - R_1^2)$  into the page (B)  $\frac{\pi I}{2}(R_1^2 - R_2^2)$  into the page  
 (C)  $\frac{\pi I}{2}(R_2^2 - R_1^2)$  out to the page (D)  $\frac{\pi I}{2}(R_2^2 + R_1^2)$  out to the page

Ans. A

Sol. As the point O is along the length of the straight wire, so the field at O due to them will be zero and hence

$$\vec{B} = \frac{\mu_0}{4\pi} \left[ \frac{\pi I}{R_2} \otimes + \frac{\pi I}{R_1} \right] \odot$$

$$\Rightarrow \vec{B} = \frac{\mu_0}{4\pi} \pi I \left[ \frac{1}{R_1} - \frac{1}{R_2} \right] \text{ out of the page}$$

$$\text{and } \vec{M} = I \left[ \frac{1}{2} \pi R_2^2 \otimes + \frac{1}{2} \pi R_1^2 \right] \odot$$

$$= \frac{1}{2} \pi I [R_2^2 - R_1^2] \text{ into the page}$$

36. The mass of the three wires of copper are in the ratio 1 : 3 : 5. and their lengths are in ratio 5 : 3 : 1 . The ratio of their electrical resistance is

(A) 1 : 3 : 5

(B) 5 : 3 : 1

(C) 1 : 15 : 125

(D) 125 : 15 : 1

Ans. D

$$\text{Sol. } R = \frac{\rho l}{A} = \frac{\rho l^2}{Al} = \frac{\rho l^2}{V} = \frac{\rho l^2}{m/d}$$

$$R = \frac{\rho d l^2}{m} \quad \text{or } R \propto \frac{l^2}{m}$$

$$R_1 : R_2 : R_3 = \frac{l_1^2}{m_1} : \frac{l_2^2}{m_2} : \frac{l_3^2}{m_3}$$

$$= \frac{25}{1} : \frac{9}{3} : \frac{1}{5} = 125 : 15 : 1$$


37. The velocities are in ground frame and the cylinder is performing pure rolling on the plank. Velocity of point 'A' would be

(A)  $2V_c$

(B)  $2V_c + V_p$

(C)  $2V_c - V_p$

(D)  $2(V_c + V_p)$



Ans. C

$$\text{Sol. } V_{Ap} = 2V_c$$





$$V_{Ag} = V_{Ap} + V_{Pg} = 2V_c - V_p$$

38. A cylinder rolls up an inclined plane, reaches some height and then rolls down (without slipping throughout these motions) The directions of the frictional force acting on the cylinder are
- (A) up the incline while ascending and down the incline while descending
  - (B) up the incline while ascending and up the incline while descending
  - (C) Down the incline while ascending and up the incline while descending.
  - (D) Down the incline while ascending as well as descending.

Ans. B

Sol. Friction force always acts up the incline irrespective of direction of motion

39. A circular platform is free to rotate on a horizontal plane about a vertical axis passing through its centre. A tortoise is sitting at the edge of the platform. Now the platform is given an angular velocity  $\omega_0$ . When the tortoise moves along a chord of the platform with a constant velocity (with respect to the platform), the angular velocity of the platform  $w(t)$  will vary with time  $t$  as



Ans. B

Sol. Conserve angular momentum

40. The image of an object, formed by a plano-convex lens at a distance of 8 m behind the lens, is real and is one-third the size of the object. The wavelength of light inside the lens is  $\frac{2}{3}$  times the wavelength in free space. The radius of the curved surface of the lens is
- (A) 1 m
  - (B) 2 m
  - (C) 3 m
  - (D) 6 m

Ans. C

Sol.  $\mu = \frac{3}{2}$  ;  $V = 8$  ;  $m = \frac{1}{3}$   
 $1 + m = \frac{V}{f} = \frac{V}{2R}$

## CHEMISTRY

41. The magnitude of an orbital angular momentum vector is  $\sqrt{6} \frac{h}{2\pi}$ . Into how many components will the vector split if a magnetic field is applied on it?

- (A) 3 (B) 5  
(C) 7 (D) 9

Ans. B

Sol. Orbital angular momentum =  $\sqrt{l(l+1)} \frac{h}{2\pi} = \sqrt{6} \frac{h}{2\pi}$

$\therefore l = 2 \rightarrow$  represents d-orbitals which splits into five components in presence of magnetic field.

42. Which of the following compound does not exist?

- (A)  $CF_4$  (B)  $SF_4$   
(C)  $OF_4$  (D)  $XeF_4$


Ans. C

Sol. Oxygen can't form more than two covalent bonds.

43. Which of the following solution mixture exerts common ion effect?

- (A)  $HCl + NaCl$  (B)  $NaHS + H_2S$   
(C)  $NaNO_3 + HNO_3$  (D)  $Na_2S + H_2SO_4$

Ans. B



44. The half-life of a chemical reaction is expressed as:  $t_{1/2} = \frac{\sqrt{3}-K}{2} C_0^{-2}$  where  $C_0$  is the initial concentration of the reactant. What is the order of the reaction?

- (A) Zero (B) 1  
(C) 2 (D) 3

Ans. D





Sol.  $t_{1/2} = \frac{\sqrt{3}-K}{2} C_0^{-2} = K C_0^{-2}$



Since  $t_{1/2} \propto C_0^{1-n}$

$$\therefore 1 - n = -2 \Rightarrow n = 3$$

45. Consider the following equilibrium constants



The correct relation among the above equilibrium constants is:

(A)  $K_4 = K_1 + K_2 + K_3$

(B)  $K_4 = K_1 \times K_2 \times K_3$

(C)  $K_4 = \frac{1}{K_1} + \frac{1}{K_2} + \frac{1}{K_3}$

(D)  $K_4 = \frac{1}{K_1} \times \frac{1}{K_2} \times \frac{1}{K_3}$

Ans. B

Sol.  $K_4 = K_1 \times K_2 \times K_3 \times K_4$   
 $= \frac{[B]}{[A]} \times \frac{[C]}{[B]} \times \frac{[D]}{[C]} = \frac{[D]}{[A]} = K_4$

46. Element 'X' belongs to the fourth period. The magnetic moment of  $X^{3+}$  ion is 5.92 B.M. Therefore, 'X' is

(A) Ni

(B) Fe

(C) Mn

(D) Co

Ans. B


Sol.  $\sqrt{n(n+2)} = 5.92$

On solving  $n = 5$

$\therefore$  The atom contains 5 unpaired electrons in +3 oxidation state

$\therefore$  The ion is  $Fe^{2+}$  and X is Fe

47.



The largest bond angle observed in the above molecule is

(A)  $\angle HCCl$

(B)  $\angle HCF$

(C)  $\angle HCH$

(D)  $\angle FCCl$

Ans. C

Sol. Carbon is more electronegative than hydrogen and F and Cl are more electronegative than carbon.

48. Which of the following pair of compounds are not isomorphous to each other?  
 (A)  $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$  and  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$  (B)  $\text{KNO}_3$  and  $\text{KClO}_3$   
 (C)  $\text{Cu}_2\text{S}$  and  $\text{Ag}_2\text{S}$  (D) None of these

Ans. B

Sol. Hybridization of N in  $\text{NO}_3^-$  is  $\text{sp}^2$  and that of Cl in  $\text{ClO}_3^-$  is  $\text{sp}^3$

49. If a blue litmus paper is dipped in  $\text{HClO}$  solution, it turns  
 (A) green (B) red  
 (C) blue (D) colourless

Ans. D

Sol.  $\text{HClO} \rightleftharpoons \text{HCl} + [\text{O}]$

Coloured substance +  $[\text{O}] \rightarrow$  Colourless substance.

50. In metallurgy Serpеч's process is used to purify  
 (A) Haematite (B) Bauxite  
 (C) Siderite (D) Calamine

Ans. B

Sol. It is used for the purification of bauxite.

51.  $\left[ \text{Fe}(\text{CO})_3(\text{PH}_3)_3 \right]$  (I)  $\left[ \text{Fe}(\text{CO})_3(\text{PH}_3)_2(\text{NH}_3) \right]$  (II)  
 $\left[ \text{Fe}(\text{CO})_3(\text{PH}_3)(\text{NH}_3)_2 \right]$  (III)  $\left[ \text{Fe}(\text{CO})_3(\text{NH}_3)_3 \right]$  (IV)


Arrange the above complexes in decreasing order of Fe – C bond energy.

- (A) I > II > III > IV (B) IV > III > II > I  
 (C) IV > II > III > I (D) I > III > IV > II

Ans. B

Sol. CO and  $\text{PH}_3$  form back bond with metal ion.

52. How many geometrical isomers are possible for the following complex?






- (A) 2 (B) 4  
 (C) 5 (D) 6

Ans. C

Sol. Five geometrical isomers are formed.

53. Which of the following electronic transition needs highest amount of energy?







(D) all the above transitions need same amount of energy

Ans. A


Sol. Orbitals with lower value of 'n' requires higher energy for electron excitation.

54. Which of the following can be easily cleaved with dil.HCl?



Ans. B

Sol. It will form the most stable carbocation.



Ammonium acetate is most soluble in


(A) Water

(B) Dil.HCl

(C) Aq.NaCl

(D) Aq.KNO<sub>3</sub>

Ans. B



∴ Forward reaction is driven up.

56. The molecular mass of a polyhydric alcohol increases by 336 amu if it reacts with  $\text{CH}_3\text{COCl}$  in presence of pyridine to form the corresponding ester. How many OH groups are present in the alcohol?


(A) 6

(B) 10

(C) 8


(D) 12

Ans. C







$\text{CH}_3\text{CO}$  (mol. mass = 43) replaces one H-atom. Therefore the mass gain by alcohol is 42 for one OH group.

$$\therefore \text{No. of OH groups} = \frac{336}{42} = 8$$




The product of above reaction is







Ans. B


Sol. Hofmann's bromamide reactions.



The end product (C) in the above reaction sequence is



Ans. A



59. How many mole of phenyl hydrazine can completely react with one mole of glucose to form glucosazone?

(A) 2

(B) 3

(C) 4

(D) 6

Ans. B

Sol. 3 moles

60. Two flasks A and B of equal volumes maintained at temperature 300 K and 700 K contain equal mass of He(g) and N<sub>2</sub>(g) respectively. What is the ratio of total translational kinetic energy of gas in flask A to that in flask B?

(A) 1:3

(B) 3:1

(C) 3:49

(D) None of these

Ans. B

Sol. 
$$\frac{\text{K.E of He}}{\text{K.E of N}_2} = \frac{\frac{3}{2}n_1RT_1}{\frac{3}{2}n_2RT_2} = \frac{\frac{3}{2} \times \frac{w}{4} \times R \times 300}{\frac{3}{2} \times \frac{w}{28} \times R \times 700} = 3:1$$

## PART – II

### MATHEMATICS

61. In the isosceles triangle  $ABC$ ,  $|\vec{AB}| = |\vec{BC}| = 8$ , a point  $E$  divides  $AB$  internally in the ratio 1 : 3, then the cosine of the angle between  $\vec{CE}$  and  $\vec{CA}$  is (where  $|\vec{CA}| = 12$ )

- (A)  $-\frac{3\sqrt{7}}{8}$  (B)  $\frac{3\sqrt{8}}{17}$   
 (C)  $\frac{3\sqrt{7}}{8}$  (D)  $\frac{-3\sqrt{8}}{17}$

Ans. C

Sol.  $\cos A = \frac{8^2 + 12^2 + 8^2}{2(8)(12)} = \frac{3}{4} \Rightarrow CE^2 = AE^2 + AC^2 - 2AE \cdot AC \cdot \cos A$


Now use cosine formula

$$|\vec{b} - \vec{c}| = |\vec{b}| = 8$$

$$b^2 + c^2 - 2\vec{b} \cdot \vec{c} = b^2 = 64$$

$$c^2 = 2\vec{b} \cdot \vec{c}$$

$$\vec{b} \cdot \vec{c} = 72$$



62. Let  $f(x) = x^3 - \frac{3x^2}{2} + x + \frac{1}{4}$ . The value of  $\int_{1/4}^{3/4} f(f(x)) dx$

- (A)  $\frac{1}{4}$  (B)  $\frac{1}{2}$   
 (C) 0 (D)  $\frac{3}{4}$

Ans. A

Sol.  $4f(x) = x^4 - (1-x)^4 + 2$   
 $\Rightarrow f(x) + f(1-x) = 1$

63. The minimum value of  $f(x) = 8^x + 8^{-x} - 4(4^x + 4^{-x}) \forall x \in \mathbb{R}$ .

- (A) -1 (B) -2  
 (C) -3 (D) 1

Ans. C

Sol. Let  $u = 2^x + 2^{-x}$

$$4^x + 4^{-x} = 4^u - 2$$

$$8^x + 8^{-x} = u^3 - 3u$$

$$\Rightarrow f(x) = u^3 - 3u - 4(u^2 - 2) = u^3 - 4u^2 - 3u + 8$$

$$\text{Let } g(u) = u^3 - 4u^2 - 3u + 8; u \geq 2$$

$$g'(u) = (3u + 1)(u - 3) \Rightarrow u = 3$$

$$g''(u) = 6u - 8 \Rightarrow g''(3) > 0 \Rightarrow u = 3 \text{ is point of minimum}$$

$$g(3) = 27 - 36 - 9 + 8 = -10$$

64. The slope of the normal at the point with abscissa  $x = -2$  of the graph of the function

$$f(x) = |x^2 - x| \text{ is}$$

(A)  $\frac{-1}{6}$

(B)  $\frac{-1}{3}$

(C)  $\frac{1}{6}$

(D)  $\frac{1}{3}$

Ans. D

Sol. For  $x < 0$

$$f(x) = |x^2 + x| = |x||x + 1|$$

For  $x < -1$

$$f(x) = (-x)(-x - 1) = x^2 + x$$

$$\therefore f'(x) = 2x + 1$$

$$\text{Slope of tangent} = 2(-2) + 1 = -3$$

$$\therefore \text{Slope of normal} = \frac{1}{3}$$

65. If  $f: \mathbb{R} \rightarrow \mathbb{R}$  is the function defined by  $f(x) = \frac{e^{x^2} - e^{-x^2}}{e^{x^2} + e^{-x^2}}$ , then

(A)  $f(x)$  is an increasing function

(B)  $f(x)$  is a decreasing function

(C)  $f(x)$  is onto (surjective)

(D) none of these

Ans. D

$$\text{Sol. } \therefore f(x) = \frac{e^{x^2} - e^{-x^2}}{e^{x^2} + e^{-x^2}} \therefore f'(x) = \frac{8x}{(e^{x^2} + e^{-x^2})^2}$$

$$= \begin{cases} > 0, & x > 0 \\ < 0, & x < 0 \\ 0, & x = 0 \end{cases}$$

66. A cylindrical gas container is closed at the top and open at the bottom; if the iron plate of the top is  $\frac{5}{4}$  times as thick as the plate forming the cylindrical sides. The ratio of the radius to the height of the cylinder using minimum material for the same capacity is
- (A)  $\frac{2}{3}$  (B)  $\frac{1}{2}$   
 (C)  $\frac{4}{5}$  (D)  $\frac{1}{3}$

Ans. C

Sol.  $V = \pi r^2 h$

If  $k$  be the thickness of the sides, then that of the top will be  $\left(\frac{5}{4}\right)k$ .

$$\therefore S = (2\pi r h)k + (\pi r^2) \left(\frac{5}{4}\right)k \quad ('S' \text{ is volume of material used})$$

$$\text{or } S = 2\pi r k \cdot \frac{V}{\pi r^2} + \frac{5}{4} \pi r^2 k = k \left( \frac{2V}{r} + \frac{5}{4} \pi r^2 \right)$$

$$\therefore \frac{ds}{dr} = k \left( -\frac{2V}{r^2} + \frac{5}{2} \pi r \right) \quad \therefore r^3 = \frac{4V}{5\pi}$$

$$\frac{d^2S}{dr^2} = k \left( \frac{4V}{r^3} + \frac{5}{2} \pi \right) = \pi k \left( 5 + \frac{5}{2} \right) = +ive$$

$$\text{when } r^3 = \frac{4V}{5\pi} \text{ or } 5\pi r^3 = 4\pi r^2 h$$

$$\therefore \frac{r}{h} = \frac{4}{5}$$

67. Let ABC be a triangle. Let A be the point (1, 2),  $y = x$  be the perpendicular bisector of AB and  $x - 2y + 1 = 0$  be the angle bisector of angle C. If the equation of BC is given by  $ax + by - 5 = 0$ , then the value of  $a + b$  is
- (A) 1 (B) 2  
 (C) 3 (D) 4


Ans. B

Sol. Image of A, say  $A'$ , w.r.t  $x - 2y + 1 = 0$

lies on BC.

$$\frac{x-1}{1} = \frac{y-2}{-2} = -2 \frac{(1-4+1)}{1+4} = \frac{4}{5}$$

$$A' = \left( \frac{9}{5}, \frac{2}{5} \right)$$





68. Let  $f(x)$  be a non – constant twice differentiable function on  $\mathbb{R}$  such that  $f(2+x) = f(2-x)$  and  $f'\left(\frac{1}{2}\right) = f'(1) = 0$ . The minimum number of roots of the equation  $f''(x) = 0$  in  $(0, 4)$  is
- (A) 2 (B) 4  
(C) 5 (D) 6

Ans. B

Sol.  $f(2+x) = f(2-x)$  and  $f'\left(\frac{1}{2}\right) = f'(1) = 0$   
It is symmetric w.r. to the line  $x = 2$ , hence  $f'(2) = 0$ .  
 $f'\left(\frac{1}{2}\right) = f'\left(2 - \frac{3}{2}\right) = -f'\left(2 + \frac{3}{2}\right)$   
 $f'(1) = f'(2-1) = -f'(2+1)$   
Thus,  $f'(x) = 0$  has roots at  $x = \frac{1}{2}, 1, 2, 3, \frac{7}{2}$ .  
 $f'(x) = 0$  has five roots  
So,  $f''(x) = 0$  will have four roots (at the least).

69. The value of  $\sum_{r=0}^{20} r(20-r) \binom{20}{r}^2$  is equal to
- (A)  $400 \times {}^{39}C_{20}$  (B)  $400 \times {}^{40}C_{19}$   
(C)  $400 \times {}^{39}C_{19}$  (D)  $400 \times {}^{38}C_{20}$

Ans. D

Sol.  $\sum_{r=0}^{20} r(20-r) \binom{20}{r}^2 = \sum_{r=0}^{20} (r \times \binom{20}{r}) ((20-r) \times \binom{20}{20-r})$   
 $= \sum_{r=0}^{20} 20 \times {}^{19}C_{r-1} \times 20 \times {}^{19}C_{19-r}$   
 $= 400 \sum_{r=0}^{20} {}^{19}C_{r-1} \times {}^{19}C_{19-r}$   
 $= 400 \times \text{coefficient of } x^{18} \text{ in } (1+x)^{19} (1+x)^{19}$   
 $= 400 \times {}^{38}C_{18}$   
 $= 400 \times {}^{38}C_{20}$

70. All the roots of the equation  $11z^{10} + 10iz^9 + 10iz - 11 = 0$  lie
- (A) inside  $|z| = 1$  (B) on  $|z| = 1$   
(C) outside  $|z| = 1$  (D) cannot say

Ans. B

Sol.  $11z^{10} + 10iz^9 + 10iz - 11 = 0$

or  $z^9(11z + 10i) = 11 - 10iz$

or  $z^9 = \frac{11 - 10iz}{11z + 10i}$

or  $|z^9| = \frac{|11i - 10z|}{|11z + 10i|}$

Now  $|11i - 10z|^2 - |11z + 10i|^2 = 21(1 - |z|)$

For  $|z| < 1$

$|11zi - 10z|^2 - |11z + 10i|^2 > 0$

$\Rightarrow |z^2| = \frac{|11i - 10z|}{|11z + 10i|} > 1$

i.e.  $|z^9| > 1$  which contradicts with  $|z| < 1$

For  $|z| > 1$  we get  $|z^9| < 1$

$\Rightarrow |z| = 1$

## PHYSICS

71. A conducting liquid bubble of radius  $a$  and thickness  $t$  ( $t \ll a$ ) is charged to potential  $V$ . If the bubble collapses to a droplet, find the potential on the droplet.

(A)  $V \left( \frac{a}{3t} \right)^{1/3}$

(B)  $Va^{1/3}$

(C)  $V \left( \frac{a^2}{t^{2/3}} \right)$

(D)  $\frac{Va^{1/3}}{3g^3}$

Ans. A

Sol.  $V = \frac{1}{4\pi\epsilon_0} \frac{q}{a}$  (for bubble)


For droplet :-  $\frac{4}{3}\pi r^3 = \frac{4}{3}\pi(a+t)^3 - \frac{4}{3}\pi a^3$

$\Rightarrow r^3 = 3a^2t \Rightarrow r = (3a^2t)^{1/3}$

$V_{\text{droplet}} = \frac{1}{4\pi\epsilon_0} \frac{q}{r} = V \left[ \frac{a}{3t} \right]^{1/3}$

72. A very long wire carrying a current 10A is bent at right angle at O. Find the magnetic induction at a point P lying on the perpendicular to the wire at O. The distance of the point P from O is 35cm.

- (A)  $0.404 \times 10^{-5}$  T  
 (B)  $0.405 \times 10^{-5}$  T  
 (C)  $0.408 \times 10^{-5}$  T  
 (D)  $0.454 \times 10^{-5}$  T



Ans. A

Sol.

$$B_1 = \frac{\mu_0 i}{4\pi R}$$

$$B_2 = \frac{\mu_0 i}{4\pi R}$$

$$B = \sqrt{B_1^2 + B_2^2}$$

$$= B_1 \sqrt{2}$$

$$= \frac{\mu_0 i \sqrt{2}}{4\pi R}$$

$$= \frac{4\pi \times 10^{-7} \times 10 \sqrt{2}}{4\pi \times 35 \times 10^{-2}}$$

$$B = 0.404 \times 10^{-5} \text{ T}$$

73. A uniform electric field of magnitude  $E = 100$  kV/m is directed upward. Perpendicular to  $E$  and directed into the page there exists a uniform magnetic field of magnitude  $B = 0.5$ T. A beam of particles of charge  $+q$  enters this region. What should be the chosen speed of particles for which the particles will not be deflected by the electric and magnetic field?

- (A)  $2 \times 10^{-5}$  m/s  
 (B)  $3 \times 10^{-5}$  m/s  
 (C)  $5 \times 10^{-5}$  m/s  
 (D)  $6 \times 10^{-5}$  m/s

Ans. A

Sol.

$$\vec{F} = q\vec{E} + q(\vec{V} \times \vec{B})$$

$$\vec{F} = q(\vec{E} + \vec{V} \times \vec{B})$$

If there has to be no deflection of beam then

$$\vec{F} = 0$$

$$\vec{E} + (\vec{V} \times \vec{B}) = 0$$

$$\vec{V} \times \vec{B} = -\vec{E}$$

$$\vec{V} \times \vec{B} = -\vec{E}$$

$$VB \sin 90^\circ = E$$

$$V = \frac{E}{B} = \frac{100 \times 10^3}{0.5}$$

$$V = 2 \times 10^5 \text{ m/sec}$$


74. In the circuit shown below, determine the current through 5V cell

(A)  $\frac{1}{15}$  Amp

(B)  $\frac{4}{15}$  Amp

(C)  $\frac{1}{17}$  Amp

(D)  $\frac{4}{17}$  Amp



Ans. A

Sol.  $\phi_A - \phi_B = 16$  ... (i)

Apply KVL on loop (cd efc)

$$0 - 5I_1 - 5 + 2(I - I_1) = 0$$

$$\Rightarrow I_1 = \frac{1}{7}(2I - 5) \quad \dots (ii)$$

Apply KVL on path (AC d B)

$$\phi_A - 2I - 5I_1 - 5 - 3I = \phi_B$$

$$\Rightarrow \phi_A - \phi_B = 5I + 5I_1 + 5$$

$$\Rightarrow 16 = 5I + 5\left(\frac{2I - 5}{7}\right) + 5$$


$$\Rightarrow 16 - 5 + \frac{25}{7} = I\left(5 + \frac{10}{7}\right)$$

$$\Rightarrow I = \frac{34}{15} \text{ Amp}$$

(i) and (ii) give

$$I_1 = -\frac{1}{15} \text{ Amp}$$

$$\text{Current through battery (5V)} = \frac{1}{15} \text{ Amp}$$



75. What amount of heat will be generated in a coil of resistance R due to a charge q passing through it if the current in the coil decreases down to zero uniformly during a time interval  $\Delta t$ ?

(A)  $\frac{4}{3} \frac{q^2 R}{\Delta t}$

(B)  $\frac{2}{3} \frac{q^2 R}{\Delta t}$

(C)  $\frac{3}{4} \frac{q^2 R}{\Delta t}$

(D)  $\frac{3}{2} \frac{q^2 R}{\Delta t}$

Ans. A

Sol. Suppose initial current is  $i_0$ , then


$$i(t) = i_0 \left(1 - \frac{t}{\Delta t}\right)$$

$$q = i_0 \int_0^{\Delta t} \left(1 - \frac{t}{\Delta t}\right) dt$$

So,  $i_0 = \frac{2q}{\Delta t}$

$$H = \int_0^{\Delta t} \left\{ \frac{2q}{\Delta t} \left( 1 - \frac{t}{\Delta t} \right) \right\}^2 R dt$$

76. A cabin is accelerating up the incline with acceleration  $g \text{ m/s}^2$ . A simple pendulum of length  $\sqrt{3}$  meter is hanging from the vertical wall of the cabin. The minimum speed given to the bob so that it performs vertical circular motion with respect to cabin is



( $g = 10 \text{ m/s}^2$ )

(A) 110 m/s

(B)  $\sqrt{15}$  m/s

(C)  $\sqrt{150}$  m/s

(D) 12 m/s

Ans. C

Sol. Apply work energy theorem and Newton's second law which will give  $u = \sqrt{150}$  m/s.

77. A particle moves along x-axis. The position of the particle at time t is given as

$$x = t^3 - 9t^2 + 24t + 1$$

The distance traveled in first 5 seconds is

(A) 20 m

(B) 10 m

(C) 18 m

(D) 28 m

Ans. D

Sol. Distance Travelled =  $\int_0^5 |\dot{x}| dt = \int_0^5 |3t^2 - 18t + 24| dt = 28$


78. A sphere of mass m has to purely roll on a rough inclined plane of coefficient of friction ' $\mu$ '. The friction force acting on the sphere is

(A)  $\mu mg \cos\theta$

(B)  $\frac{2mg \sin\theta}{7}$  downward

(C)  $\frac{2mg \sin\theta}{7}$  upward

(D)  $\frac{5mg \sin\theta}{7}$  downward



Ans. C

Sol.  $f = \frac{mg \sin\theta}{1 + \frac{R^2}{K^2}}$

79. A body of mass 1kg is suspended from a massless spring having force constant 600 N/m. Another body of mass 0.5 kg moving vertically upwards hits the suspended body with a velocity 3 m/sec and gets embedded in it. The frequency of oscillation and the amplitude of motion are

- (A)  $\frac{5}{\pi}$  Hz, 10Cm  
 (B)  $\frac{10}{\pi}$  Hz, 5Cm  
 (C)  $\frac{5}{\pi}$  Hz, 5Cm  
 (D)  $\frac{5}{\pi}$  Hz, 10Cm

Ans. B

Sol.  $V = \frac{0.5 \times 3}{(0.5 + 1)} = 1 \text{ m/s} \Rightarrow K_{\max} = U_{\max}$

$$\frac{1}{2} \times 1.5 \times 1^2 = \frac{1}{2} \times 600 \times A^2$$

A = 5 cm

$$f = \frac{1}{2\pi} \sqrt{\frac{k}{m}} = \frac{1}{2\pi} \sqrt{\frac{600}{1.5}} = \frac{10}{\pi}$$

$$f = \frac{10}{\pi} \text{ Hz}$$

80. When the electron in a hydrogen atom jumps from the second orbit to the first orbit, the wavelength of the radiation emitted is  $\lambda$ . When the electron jumps from the third to the first orbit, the wavelength of the radiation emitted as

- (A)  $\frac{9}{4}\lambda$   
 (B)  $\frac{4}{9}\lambda$   
 (C)  $\frac{27}{32}\lambda$   
 (D)  $\frac{32}{27}\lambda$

Ans. C


Sol.  $\frac{hc}{\lambda} = R \left( \frac{1}{1^2} - \frac{1}{2^2} \right) = \frac{3}{4}R \quad \dots(i)$

$$\frac{hc}{\lambda'} = R \left( \frac{1}{1^2} - \frac{1}{3^2} \right) = \frac{8}{9}R \quad \dots(ii)$$

$$\frac{\lambda'}{\lambda} = \frac{3}{4} \times \frac{9}{8}, \quad \lambda' = \frac{27}{32}\lambda$$

## CHEMISTRY

81. The average energy of each hydrogen bond in A-T pair is  $x \text{ kcal mol}^{-1}$  and that in G-C pair is  $y \text{ kcal mol}^{-1}$ . Assuming that no other interaction exists between the nucleotides, the approximate energy required in  $\text{kcal mol}^{-1}$  to split the following double stranded DNA into two single strands is







[Each dashed line may represent more than one hydrogen bond between the base pairs]

- (A)  $10x + 9y$                       (B)  $5x + 3y$   
(C)  $15x + 6y$                       (D)  $5x + 4.5y$

Ans. A


- Sol. Number of H-bond is A-T pair = 2, while no of H-bond in G-C pair is 3. Therefore  
(i) Total number of A-T. H-bond = number of A-T pair  $\times$  Number of H bond =  $5 \times 2 = 10$   
(ii) Total number of G-C H-bond = number of G-C pair  $\times$  number of H bond =  $3 \times 3 = 9$   
Total energy required to dissociate the stand =  $10x + 9y \text{ Kcal mol}^{-1}$

82. The correct representation of wavelength intensity relationship of an ideal blackbody radiation at two different temperatures  $T_1$  and  $T_2$  is



Ans. A

Sol.




83. A gas at atmospheric pressure is heated from 0°C to 546°C and simultaneously compressed to one-third of its original volume. Hence final pressure is.  
 (A) 6 atm (B) 9 atm  
 (C) 18 atm (D) 27 atm

Ans. B

Sol.  $P_1 = 1 \text{ atm}$   
 $T_1 = 273 \text{ K}$   
 Let volume at this condition is  $V \text{ L}$  and  
 $P_2 = P \text{ atm}$   
 $T_2 = 819 \text{ K}$   
 $V_2 = \frac{V}{3} \text{ L}$   
 Then,  $\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$   
 $\Rightarrow \frac{1 \times V}{273} = \frac{P \times V}{3 \times 819}$   
 $\Rightarrow P = 9 \text{ atm}$   
 Hence, the correct answer is option B.

84. Alkanamide which on Hoffmann's reaction gives 1-phenyl ethyl amine is:  
 (A) 2-phenylpropanamide (B) 3-phenylpropanamide  
 (C) 2-phenylethanamide (D) N-phenylethanamide

Ans. A



Hence, the correct answer is option A.

85. The order of  $pK_a$  of these carboxylic acids in water is  
 (I)  $\text{CH}_3\text{COOH}$  (II)  $\text{C}_2\text{H}_5\text{COOH}$   
 (III)  $\text{C}_3\text{H}_7\text{COOH}$   
 (A)  $\text{I} > \text{II} > \text{III}$  (B)  $\text{II} > \text{I} > \text{III}$   
 (C)  $\text{III} > \text{II} > \text{I}$  (D)  $\text{III} > \text{I} > \text{II}$

Ans. C

Sol. The electron releasing substituents (+I) decrease the acidic strength of carboxylic acid by destabilizing the carboxylate ion. The strength of the acid is expressed in terms of the dissociation constant ( $K_a$ ). A stronger acid has higher  $K_a$  but lesser  $pK_a$  value. Hence, the correct answer is option C.



86. X mL of O<sub>2</sub> effuses through a hole in a container in 20 seconds. The time taken for the effusion of the same volume of the gas specified below under identical conditions is  
 (A) 10 seconds : He (B) 5 seconds : H<sub>2</sub>  
 (C) 25 seconds : CO (D) 55 seconds : CO<sub>2</sub>

Ans. B

Sol.  $\frac{r_{\text{Gas}}}{r_{\text{O}_2}} = \sqrt{\frac{M_{\text{O}_2}}{M_{\text{Gas}}}}$ , where r is rate of effusion and M is molecular mass of gas.

$$\Rightarrow \frac{V_{\text{Gas}}}{t_{\text{Gas}}} \times \frac{t_{\text{O}_2}}{V_{\text{O}_2}} = \sqrt{\frac{M_{\text{O}_2}}{M_{\text{Gas}}}}, \text{ where t is time for effusion of V is volume of gas effused.}$$

If  $V_{\text{O}_2} = V_{\text{Gas}} = X \text{ mL}$

$$\Rightarrow \frac{t_{\text{O}_2}}{t_{\text{Gas}}} = \sqrt{\frac{M_{\text{O}_2}}{M_{\text{Gas}}}}$$


$$\frac{20}{t_{\text{Gas}}} = \sqrt{\frac{32}{M_{\text{Gas}}}}$$

Hence, the correct answer is option B.

87. How many Faradays of electricity is needed to oxidise one mole of H<sub>2</sub>O completely to dioxygen gas?  
 (A) 1 (B) 2  
 (C) 4 (D) 1.5

Ans. B

Sol. One mole H<sub>2</sub>O contains one mole O<sup>2-</sup>



88. Which of the following atom has most stable electronic configuration in +2 oxidation state? [At. No. of Cr = 24, Mn = 25, Fe = 26, Cu = 29]  
 (A) Cr (B) Mn  
 (C) Fe (D) Cu


Ans. B

Sol. Mn(4s<sup>2</sup>3d<sup>5</sup>)





Mn<sup>2+</sup> contains the stable 3d<sup>5</sup> half-filled electron configuration.

89. Which of the following releases SO<sub>2</sub> gas when treated with dil.H<sub>2</sub>SO<sub>4</sub>?  
 (A) Na<sub>2</sub>SO<sub>4</sub> (B) Na<sub>2</sub>SO<sub>3</sub>  
 (C) Na<sub>2</sub>S (D) All are correct

Ans. B



90. Which of the following complex displays ionization isomerism?



Ans. D

Sol. In(D) the  $\text{Br}^-$  and  $\text{Cl}^-$  ligands exchange their positions.