



Resonance
Educating for better tomorrow

KISHORE VAIGYANIK PRO TSAHAN YOJANA - 2014

Date : 02-11-2014

Duration : 3 Hours

Max. Marks : 100

STREAM - SA

GENERAL INSTRUCTIONS

- The Test Booklet consists of **80** questions.
- There are Two parts in the question paper. The distribution of marks subjectwise in each part is as under for each correct response.

MARKING SCHEME :

PART-I :

MATHEMATICS

Question No. **1 to 15** consist of **ONE (1)** mark for each correct response.

PHYSICS

Question No. **16 to 30** consist of **ONE (1)** mark for each correct response.

CHEMISTRY

Question No. **31 to 45** consist of **ONE (1)** mark for each correct response.

BIOLOGY

Question No. **46 to 60** consist of **ONE (1)** mark for each correct response.

PART-II :

MATHEMATICS

Question No. **61 to 65** consist of **TWO (2)** marks for each correct response.

PHYSICS

Question No. **66 to 70** consist of **TWO (2)** marks for each correct response.

CHEMISTRY

Question No. **71 to 75** consist of **TWO (2)** marks for each correct response.

BIOLOGY

Question No. **76 to 80** consist of **TWO (2)** marks for each correct response.



Resonance Eduventures Pvt. Ltd.

CORPORATE OFFICE : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005
Tel. No. : 0744-3192222, 3012222, 3022222 | Toll Free : 1800 200 2244 | To Know more : sms RESO at 56677
Website : www.resonance.ac.in | Email : contact@resonance.ac.in

This solution was download from Resonance KVPY 2014 Solution portal

PART-I

One Mark Questions

MATHEMATICS

1.

Sol. r be a root $\Rightarrow r^2 + 2r + 6 = 0$ (1)

$$\begin{aligned} &\text{now } (r+2)(r+3)(r+4)(r+5) \\ &= (r^2 + 5r + 6)(r^2 + 9r + 20) \\ &= (3r)(7r + 14) \quad \text{using (i)} \\ &= 21(r^2 + 2r) \\ &= -126 \quad \text{using (i)} \end{aligned}$$

Ans. (C)

2.

Sol. Given $f(x) + (x + \frac{1}{2})f(1-x) = 1$ (1)

but $x = 0$

$$f(0) + \frac{1}{2}f(1) = 1$$

$$\Rightarrow 2f(0) + f(1) = 2 \quad \text{.....(2)}$$

put $x = 1$ in (1)

$$\Rightarrow f(1) + \frac{3}{2}f(0) = 1$$

$$\Rightarrow 2f(1) + 3f(0) = 2 \quad \text{.....(3)}$$

Solving (2) & (3) we have

$$f(0) = 2 \text{ \& } f(1) = -2$$

$$\therefore 2f(0) + f(1) = 4 - 2 = 2$$

Ans. (C)

3.

$$\frac{1^3 + 2^3 + \dots + (2n)^3}{1^2 + 2^2 + \dots + n^2} = \left(\frac{2n(2n+1)}{2} \right)^2 \cdot \frac{6}{n(n+1)(2n+1)}$$

$$= \frac{6n(2n+1)}{n+1}$$

$$= \frac{12n^2 + 6n}{n+1} = \frac{12(n^2 - 1) + 6(n+1) + 6}{n+1}$$

$$= 1 + \frac{6}{n+1}$$

If the given terms is an integer, then $\frac{6}{n+1}$ must be an integer

$$\Rightarrow n = 1, 2, 5$$

$$\text{Sum} = 8$$

Ans. (A)

4.

$$X \rightarrow ab \text{ or } x = 10a + b$$

$$y \rightarrow ba \text{ or } y = 10b + a$$

$$\text{Now } x^2 - y^2 = (10a + b)^2 - (10b + a)^2$$

$$= 99(a^2 - b^2)$$

$$= 3^2 \times 11(a+b)(a-b) \quad \text{----- (1)}$$

According to Q

$$(a+b)(a-b) = 11 \text{ and } a-b = 1$$

$$\Rightarrow a+b = 11 \text{ and } a-b = 1$$

$\Rightarrow a = 6, b = 5$

Hence

$x = 65$

$y = 56$

and $m = 33 \Rightarrow x + y + m = 154$

Ans. (D)

5.

Sol.

$\therefore \text{HCF} = x - 1$

$\Rightarrow p(x) = x^2 - 5x + a$

$= x^2 - 5x + 4$

$= (x - 1)(x - 4) \dots\dots\dots(1)$

and

$q(x) x^2 - 3x + b = x^2 - 3x + 2$

$= (x - 1)(x - 2) \dots\dots\dots(2)$

$\Rightarrow k(x) = (x - 1)(x - 2)(x - 4)$

Hence

$(x - 1) + R(x) = (x - 1) + (x - 1)(x - 2)(x - 2)(x - 4)$

$= (x - 1)(x - 3)^2$

Hence sum of roots = 7


Ans. (D)

6.

Ans. (D)

7.

Sol.



area of sector $OACB = \frac{r^2}{2} \theta = \frac{1}{2} \cdot \frac{\pi}{3} = \frac{\pi}{6}$

area of shaded region = $\frac{\pi}{6} - \text{area of } \triangle OAB$

$= \frac{\pi}{6} - \frac{\sqrt{3}}{4}$

Hence area of line = Area of semi-circle – area of shaded region

$= \frac{1}{2} \pi \left(\frac{1}{2}\right)^2 - \left(\frac{\pi}{6} - \frac{\sqrt{3}}{4}\right)$

$= \frac{\sqrt{3}}{4} + \frac{\pi}{8} - \frac{\pi}{6}$

$= \frac{\sqrt{3}}{4} - \frac{\pi}{24}$

Ans. (B)

8.

Sol.

$\therefore \frac{AI}{IF} = \frac{b+c}{a} \dots\dots\dots (1)$

$\therefore \frac{BI}{ID} = \frac{a+c}{b} = \frac{3}{2} \dots\dots\dots(2)$

$$\therefore \frac{CI}{IE} = \frac{a+c}{c} = \frac{2}{1}$$

$$\Rightarrow a + b = 2c \quad \dots\dots(3)$$

$$(2) \ 2a + 2c = 3b \quad \text{using to}$$

$$\Rightarrow 2a + a + b = 3b \quad \text{using (3)}$$

$$\Rightarrow 3a = 2b$$

$$\Rightarrow b = \frac{3}{2}a \quad \dots\dots(4)$$


$$\text{Now again (3)} \Rightarrow 2c = a + b$$

$$= a + \frac{3}{2}a$$

$$\Rightarrow c + \frac{5}{4}a$$


$$\text{Hence } \frac{AI}{IF} = \frac{b+c}{a} = \frac{\frac{1}{2}a + \frac{5}{4}a}{a} = \frac{11}{4}$$

Ans. (B)



9.

Sol.



$$\therefore RP = RA = 10 - \alpha$$

$$\Rightarrow RS = 10 - \alpha + \beta \quad \dots\dots (1)$$

$$\text{Also } SQ = SD = 10 - \beta$$


$$\Rightarrow RS = 10 - \beta + \alpha \quad \dots\dots(2)$$

$$(1) \text{ and } (2) \Rightarrow \alpha = \beta, \text{ Hence } RS = 10$$

Ans. (C)

10.

Sol.



1. $\triangle AOB$ is equilateral ($\angle AOB = \angle OAB = \angle OBA = 60^\circ$)
2. $\triangle OBC$ is right angled isosceles ($\angle OBC = 90^\circ$)
3. $\triangle ABC$ is isosceles ($\angle BAC = \angle BCA = 15^\circ$)
4. $\angle OAC = 60^\circ - \angle CAB = 45^\circ$

5. $\triangle AOF$ is right angled isosceles ($\angle AOF = 90^\circ, \angle OFA = 45^\circ$)
 6. $\angle BOF = 90^\circ - \angle AOB = 30^\circ$
 7. $\triangle OBC$ is right angled isosceles ($\angle BOC = 45^\circ$)
 $\therefore \frac{\angle BOF}{\angle BOC} = \frac{30^\circ}{45^\circ} = \frac{2}{3}$

Ans. (B)**11.****Sol.**

Let total seats = 100
 on first day,
 Ticket price = 200
 seats full = 60%

$$= \frac{60}{100} \times 100 = 60$$

$$\therefore \text{Revenue} = 60 \times 200$$

$$R_1 = 12000$$

On second day

Tricked price = 200 - 20% of 200

$$= 200 - \frac{20}{100} \times 200$$

$$= 200 - 40 = 160$$

Seats full 60 + 50% of 60

$$= 60 + \frac{50}{100} \times 60$$

$$= 60 + 30 = 90$$

Revenue = 160 × 90

$$R_2 = 14400$$

$$\% \text{ Increase in Revenue} = \frac{R_2 - R_1}{R_1} \times 100$$

$$= \frac{14400 - 12000}{12000} \times 100$$

$$= \frac{2400}{1200} \times 100$$

$$= 20\%$$

Ans. (D)**12.****Sol.**

year	Population
2010	— 39
2011	— 60
2012	— x
2013	— 123

According to Q

2010 — 39

2011 — 60

2012 — x

2013 — 123

According to Q

$$x - 39 = k(60) \text{ \& } 63 = kr$$

$$\Rightarrow x - 39 = \frac{63}{x} \cdot 63$$

$$\Rightarrow x^2 - 39x = -(60)(63) = 0$$

$$x = 84 \text{ \& } -40$$

Ans(B)

13. $N = ab\ ab\ ab$
 $1 < a \leq 9 \quad 0 < b \leq 9 \quad a, b \in \mathbb{I}$
 $N = 10^5a + 10^4b + 10^3a + 10^2b + 10a + b$
 $= (10^4 + 10^2 + 1)(10a + b)$
 $= (10^2 + 10 + 1)(10^2 - 10 + 1)(10a + b)$
 $= 3 \times 37 \times 13 \times 7(10a + b) \quad \dots\dots\dots (1)$

then $10a + b = P_1 \times P_2 \quad p_1, p_2 \in \text{prime and } 10 \leq 10a + b \leq 99$

a	b	$10a + b$
1	0	$10 = 2 \times 5$
2	2	$22 = 2 \times 11$
3	4	$34 = 2 \times 17$
3	8	$38 = 2 \times 19$
4	6	$46 = 2 \times 23$
5	5	$55 = 5 \times 11$
5	8	$58 = 2 \times 29$
6	2	$62 = 2 \times 31$
7	4	$74 = 2 \times 37$
8	2	$82 = 2 \times 41$
8	5	$85 = 5 \times 17$
9	4	$94 = 2 \times 47$
9	5	$95 = 5 \times 19$

Ans(C)

14. **Sol.** Let house no are $\alpha, \alpha + 2, \alpha + 4, \alpha + 6, \alpha + 8, \alpha + 10, \dots$

$\alpha + 10 = a \Rightarrow \alpha = a - 10 \quad \dots\dots\dots (1)$
 House no. will be (+)
 $\Rightarrow \alpha = a - 10 > 0$
 $\Rightarrow \alpha > 10$
 $\Rightarrow \alpha \geq 12$ as a is each too $\dots\dots\dots(2)$

Now $S_n = \frac{n}{2}[2\alpha + (n-1)d]$

$170 = \frac{n}{2}[2\alpha + (n-1)(2)]$
 $= n(\alpha + (n-1))$
 $= n(a - 10 + n - 1)$
 $= n(a - 11 + n)$
 $\Rightarrow n^2 + n(a - 11) - 170 = 0$
 $\Rightarrow n = \frac{(11-a) \pm \sqrt{(a-11)^2 + 680}}{2} \quad \dots\dots\dots(3)$

$\therefore n \geq 6$
 $\Rightarrow \frac{(11-a) \pm \sqrt{(a-11)^2 + 680}}{2} \geq 6$

$\Rightarrow a \leq \frac{800}{24} \quad \dots\dots\dots(4)$

From (2) and (4) $\Rightarrow 12 \leq a \leq 32$
 Now checking through (3) for $a = 12, 14, \dots$;
 we have $a = 18, n = 10$ and $S_n = 170$
 Hence options

Ans(C)

15.

Sol. $\frac{5}{7} = \frac{2520a_2 + 840a_3 + 210a_4 + 42a_5 + 7a_6 + a_7}{7}$

$$2520a_2 + 840a_3 + 210a_4 + 42a_5 + 7a_6 + a_7 = 3600$$

Let $a_2 = a_3 = a_4 = 1$ $a_5 = 0$ $a_6 = 4$ $a_7 = 2$

Ans(B)

PHYSICS

16.

Sol. |slope| is increasing at point R

Ans. (A)


17.

Sol. No Buoyancy force in vacuum

Ans. (D)

18.

Sol.



$$\tan \theta = \frac{F}{mg} \quad (F \rightarrow \text{same})$$

$$\tan \theta \propto \frac{1}{m}$$

$$\therefore m_1 = m_2$$

Ans. (B)


19.

Sol. Case-1

$$v = \sqrt{2gh}$$

Case-2

$$\Delta U + \Delta KE = w_f$$



$$-mgh + \frac{1}{2} m \left(\frac{2gh}{9} \right) = -\mu mgh$$

$$\mu = \frac{8}{9}$$


Ans. (A)

20. **Ans. (C)**

21. **Ans. (D)**


22.

Sol. For min deviation
 $i = e$



$$r_1 = r_2 = \frac{A}{2}$$

$$\therefore r_1 = r_2 = 30^\circ$$




For minimum deviation i should lie between 40° to 50°

Ans. (B)

23.

Sol.



$$\frac{1}{F} = \left(\frac{1.6}{2} - 1 \right) \left(\frac{1}{-0.2} - \frac{1}{0.2} \right)$$

$$= \frac{0.4}{2} \times \frac{1}{0.1}$$

$F = 0.5$ converging lens


Ans. (D)

24. **Sol.** In option B it will not move, in option C & D path will be straight line.

Ans. (A)

25.

Sol. $\mu_i = \frac{kQ^2}{d} = E$




$$\mu_f = \frac{kQ^2}{d} + \frac{k(-Q)^2}{d} + \frac{k-Q^2}{d}$$

$$= -\frac{kQ^2}{d} = -E$$

Ans. (B)

26. **Sol.** Using Lenz's law upper face first becomes North pole then south pole



Ans. (C)

27. **Ans. (B)**

28. In SHM particle comes 2 times at every position in 1 oscillation, so actual histogram may be option (A) but since at it random snap shots so it should be option (C)

Ans. (C)


29. **Ans. (B)**

30. **Ans. (A)**

CHEMISTRY

- 31.


Sol. CO & N₂ are isoelectronic



Ans. (A)

- 32.

Sol. HYDRAZINE N₂H₄



LP = 2

BP = 5

Ans. (B)

- 33.

Sol. $C(s) + O_2(g) \longrightarrow CO_2(g)$

moles = 1mole 1mole 1mole

weight = 12gm 32gm 44gm

12gm of C require \rightarrow 1 mole of O₂

\therefore 2.4gm of C will require $\rightarrow \frac{1}{12} \times 2.4$ mole of O₂

volume of 2.4/12 mole O₂ at STP = $\frac{22.4 \times 2.4}{12}$ litre

4.48 litre

Ans. (D)

34.

Sol. Nonpolar substance will have high R_f value as solvent is nonpolar therefore option (A) will have high R_f value as it have low dipole moment.

Ans. (A)35. **Ans. (A)**

36.

Sol.

$$r_n = \frac{R_H n^2}{Z}$$

$$r_{H^{e+}} = \frac{53 n^2}{Z}$$

$$= \frac{53 \times 1^2}{2} = 27 \text{ approx.}$$

Ans. (C)37. **Ans. (D)**

38.

Sol. $NH_4Cl \rightarrow$ acidic Salt ($PH < 7$) $NaCl \rightarrow$ Neutral Salt ($PH = 7$) $CH_3COONa \rightarrow$ Basic salt ($PH > 7$)**Ans. (B)**

39.

Sol. average speed $\propto \frac{1}{\sqrt{M}}$

$$\frac{V_{He}}{V_{O_2}} = \sqrt{\frac{32}{4}} = \sqrt{\frac{M_{O_2}}{M_{He}}}$$


$$= \sqrt{8} = 2\sqrt{2}$$

Ans. (A)


40.

Sol. $NH_4Cl + NaNO_2 \longrightarrow NaCl + N_2 + 2H_2O$ **Ans. (C)**

41.

Sol. $CH_3 - O - CH_2 - CH_2 - CH_3$ $CH_3 - CH_2 - O - CH_2 - CH_3$ **Ans. (B)**

42.



Oxidation

Ans. (D)

43.

Sol. I, II & IV compound form H-bond III do not form H-Bond

Ans. (C)

44.

Sol. $\Delta G^\circ = -RT \ln K_{eq}$

Ans. (C)

45.

Sol. As we move from left to right in period ionisation energy increases.

Ans. (A)

BIOLOGY

46. (B) 47. (A) 48. (B) 49. (D) 50. (B) 51. (D) 52. (C)
 53. (A) 54. (B) 55. (A) 56. (D) 57. (B) 58. (B) 59. (C)
 60. (D)

PART-II Two Mark Questions

MATHEMATICS

61. $a + b + c = 0, \quad a, b, c \in R \neq 0$
 $a^2 + b^2 + c^2 + 2(ab + bc + ca) = 0$
 $q = a^2 + b^2 + c^2, \quad r = a^4 + b^4 + c^4$
 $r = q^2 - 2(a^2b^2 + b^2c^2 + c^2a^2)$
 $r = q^2 - 2[(ab + bc + ca)^2 - 2abc(a + b + c)]$
 $r = q^2 - 2(q^2 / 4)$
 $r = q^2 / 2$

ANS - B

62. $\frac{1}{1 + \sqrt{2^{1947}}} + \frac{1}{2^{1947} + 2^{\frac{1947}{2}}} = \frac{1}{2^{\frac{1947}{2}}}$

Similarly & \therefore

$$\sum_{n=0}^{1947} \frac{1}{2^n + \sqrt{2^{1947}}} = \frac{974}{\sqrt{2^{1947}}} = \frac{487}{\sqrt{2^{1945}}}$$

ANS - A

63. $\frac{x^2 - 1 + 1}{x - 1} + \frac{y^2 - 1 + 1}{y - 1} = 4$

$$x + 1 + \frac{1}{x - 1} + y + 1 + \frac{1}{y - 1} = 4$$

$$a + 2 + \frac{1}{x-1} + \frac{1}{(a-1)-x} = 4$$

$$\frac{(a-1)-x+x-1}{(x-1)[(a-1)-x]} = 2-a$$

∴ a ≠ 2 [for a = 2 equation have infinitely many solution]

$$\therefore (x-1)[(a-1)-x] = -1$$

$$(x-1)[x-(a-1)] = 1$$

$$x^2 - ax + (a-2) = 0$$

$$D > 0$$


∴ equation have 2 real roots so

a can be 1, 3, 4,..... 2014

ans 2013

ANS - C

64.



Equation of line AB is

$$\frac{x}{\sqrt{7}} + \frac{y}{\sqrt{5}} = 1$$

$$\text{Let } P \left[\alpha, \sqrt{5} \left(1 - \frac{\alpha}{\sqrt{7}} \right) \right]$$

on solving $16(PA)^2 = 9(PB)^2$

$$P \left[\frac{\sqrt{7}}{3}, \frac{2\sqrt{5}}{3} \right]$$

$$\text{Let } BP : PC = \lambda : 1$$

$$\text{then } \lambda = 2$$

$$BP : PC = 2 : 1$$

ANS - (A)

65. $(a \times b \times c) + (a \times b) + (c \times a) + (a + b + c) = 29$

$$(1+a)(1+b)(1+c) = 30$$

$$= 2 \times 3 \times 5 \rightarrow (a, b, c) \Rightarrow (1, 2, 3) \Rightarrow 6$$

$$= 1 \times 6 \times 5 \rightarrow (a, b, c) \Rightarrow (0, 5, 4) \Rightarrow 4$$

$$= 1 \times 3 \times 10 \rightarrow (a, b, 1) \Rightarrow (0, 2, 9) \Rightarrow 4$$

14

ANS - (C)


PHYSICS

66.

Sol. Finally com at p

$$X_{\text{am}} = \frac{A_1 X_1 + A_2 X_2}{A_1 + A_2}$$

$$(a-b) = \frac{a(a-b)\frac{(a-b)}{2} + b(a-b)(a-b+b/2)}{a(a-b) + (a-b)b}$$



$$\therefore \left(\frac{a}{b}\right)^2 - \left(\frac{a}{b}\right) - 1 = 0$$

$$\frac{a}{b} = \frac{1 + \sqrt{5}}{2}$$


Ans. (B)

67. Weight = F_0
 $4\pi r^2 t \rho_w g + 4/3 \pi r^3 \rho_{Ne} g = 4/3 \pi r^3 \rho_{air} g$
 $\therefore t = 3.5 \mu m$

Ans. (D)

68. **Sol.** Heat lost = heat gas
 $0.05 \times 900 \times (300 - 160) = 1 \times 4200 \times (T - 30)$
 $T = 31.5^\circ$
Ans. (C)

69. **Sol.**




- also
 $45 + r > C$
 $45 - r > C$
 $90 > C$

$$\therefore \mu > \sqrt{2}$$

Ans (A)

70.



Using KCL

At point A

Current is 3mA

At point C

Current is 8 mA

At point E

Current is 21 mA

At point G

Current through GH is


34 ma

$$\therefore V_{PQ} = V_{GH} = i R_{GH}$$

$$= 34 \text{ V}$$

Ans. (D)**CHEMISTRY**

71.




moles of gas remaining = 9.7
 at constant (T) & (V)

$$\frac{n_1}{n_2} = \frac{p_1}{p_2}$$

$$\frac{10}{9.7} = \frac{1}{p_2} \quad \& \quad p_2 = 0.97$$

Ans. (B)

72.



$$\text{millimole of } H_2SO_4 = \frac{\text{mmol of } NH_3}{2} = 20$$

$$\text{mmol } NH_3 = \text{mmol of } N = 40$$

$$W_N = \frac{40 \times 14}{1000} = \frac{560}{1000} = 0.56g$$

$$\% \text{ of } N = \frac{0.56}{2} \times 100 = 28$$

Ans. (A)

73.

Sol. 1.125L of H_2 produced by 0.1 equivalent of metal

1.85L of H_2 will be produced by $= \frac{0.1 \times 1.85}{1.125}$ equivalents

\therefore No of gram equivalent of metal

$$= \frac{2}{\text{Equivalent weight}} = \frac{2}{x}$$

$$\therefore \frac{0.1}{1.125} \times 1.85 = \frac{2}{x}$$


$$\boxed{x = 12.16}$$

Ans. (D)

74.

Sol. $CaO + C \longrightarrow CaC_2 + CO_2$

$CaC_2 + H_2O \longrightarrow HC \equiv CH + Ca(OH)_2$



Ans. (A)

75.

Sol.

Ans. (A)

BIOLOGY

76. (C) 77. (C) 78. (B) 79. (A) 80. (A)