


5. Particle A of mass $m_A = \frac{m}{2}$ moving along the x-axis with velocity v_0 collides elastically with another particle B at rest having mass $m_B = \frac{m}{3}$. If both particles move along the x-axis after the collision, the change $\Delta\lambda$ in de-Broglie wavelength of particle A, in terms of its de-Broglie wavelength (λ_0) before collision is:

- (A) $\Delta\lambda = \frac{3}{2}\lambda_0$ (B) $\Delta\lambda = \frac{5}{2}\lambda_0$
 (C) $\Delta\lambda = 2\lambda_0$ (D) $\Delta\lambda = 4\lambda_0$

6. A wire A, bent in the shape of an arc of a circle, carrying a current of 2 A and having radius 2 cm and another wire B, also bent in the shape of arc of a circle, carrying a current of 3 A and having radius of 4 cm, are placed as shown in the figure. The ratio of the magnetic fields due to the wires A and B at the common centre O is:




- (A) 6 : 5 (B) 2 : 5
 (C) 6 : 4 (D) 4 : 6

7. The specific heat of water = $4200 \text{ J kg}^{-1}\text{K}^{-1}$ and the latent heat of ice = $3.4 \times 10^5 \text{ J kg}^{-1}$. 100 grams of ice at 0°C is placed in 200 g of water at 25°C . The amount of ice that will melt as the temperature of water reaches 0°C is close to (in grams):

- (A) 64.6 (B) 61.7
 (C) 69.3 (D) 63.8

8. Take the breakdown voltage of the zener diode used in the given circuit as 6V. For the input voltage shown in figure below, the time variation of the output voltage is: (Graphs drawn are schematic and not to scale)




- (A) (B)
 (C) (D)

9. A beam of plane polarized light of large cross-sectional area and uniform intensity of 3.3 Wm^{-2} falls normally on a polarizer (cross sectional area $3 \times 10^{-4} \text{ m}^2$) which rotates about its axis with an angular speed of 31.4 rad/s . The energy of light passing through the polariser per revolution, is close to:

- (A) $1.0 \times 10^{-5} \text{ J}$ (B) $1.5 \times 10^{-4} \text{ J}$
 (C) $5.0 \times 10^{-4} \text{ J}$ (D) $1.0 \times 10^{-4} \text{ J}$

10. A small bar magnet placed with its axis at 30° with an external field of 0.06 T experiences a torque of 0.018 Nm. The minimum work required to rotate it from its stable to unstable equilibrium position is:
 (A) 11.7×10^{-3} J (B) 9.2×10^{-3} J
 (C) 7.2×10^{-2} J (D) 6.4×10^{-2} J

11. A two point charges $4q$ and $-q$ are fixed on the x-axis at $x = -\frac{d}{2}$ and $x = \frac{d}{2}$, respectively. If a third point charge 'q' is taken from the origin to $x = d$ along the semicircle as shown in the figure, the energy of the charge will:



- (A) decrease by $\frac{4q^2}{3\pi\epsilon_0 d}$ (B) increase by $\frac{2q^2}{3\pi\epsilon_0 d}$
 (C) increase by $\frac{3q^2}{4\pi\epsilon_0 d}$ (D) decrease by $\frac{q^2}{4\pi\epsilon_0 d}$
12. Choose the correct option relating wavelengths of different parts of electromagnetic wave spectrum:
 (A) $\lambda_{x\text{-rays}} < \lambda_{\text{micro waves}} < \lambda_{\text{radio waves}} < \lambda_{\text{visible}}$ (B) $\lambda_{\text{visible}} < \lambda_{\text{micro waves}} < \lambda_{\text{radio waves}} < \lambda_{x\text{-rays}}$
 (C) $\lambda_{\text{visible}} > \lambda_{x\text{-rays}} > \lambda_{\text{radio waves}} > \lambda_{\text{micro waves}}$ (D) $\lambda_{\text{radio waves}} > \lambda_{\text{micro waves}} > \lambda_{\text{visible}} > \lambda_{x\text{-rays}}$


13. Match the C_P/C_V ratio for ideal gases with different type of molecules:

Molecule Type			C_P/C_V
(A)	Monatomic	(I)	$\frac{7}{5}$
(B)	Diatomic rigid molecules	(II)	$\frac{9}{7}$
(C)	Diatomic non-rigid molecules	(III)	$\frac{4}{3}$
(D)	Triatomic rigid molecules	(IV)	$\frac{5}{3}$

Match the correct option?

- (A) A \rightarrow III; B \rightarrow IV; C \rightarrow II; D \rightarrow I
 (B) A \rightarrow IV; B \rightarrow I; C \rightarrow II; D \rightarrow III
 (C) A \rightarrow II; B \rightarrow III; C \rightarrow I; D \rightarrow IV
 (D) A \rightarrow IV; B \rightarrow II; C \rightarrow I; D \rightarrow III

14. A small bar magnet is moved through a coil at constant speed from one end to the other. Which of the following series of observations will be seen on the galvanometer G attached across the coil?



Three positions shown describe:
 (a) the magnet's entry
 (b) magnet is completely inside and
 (c) magnet's exit

(A)

(B)

(C)

(D)

15. For a transverse wave traveling along a straight line, the distance between two peaks (crests) is 5 m, while the distance between one crest and one trough is 1.5 m. The possible wavelengths (in m) of the waves are:

- (A) 1, 3, 5, (B) $\frac{1}{1}, \frac{1}{3}, \frac{1}{5}, \dots$
 (C) $\frac{1}{2}, \frac{1}{4}, \frac{1}{6}, \dots$ (D) 1, 2, 3,


16. A air bubble of radius 1 cm in water has an upward acceleration 9.8 cm s^{-2} . The density of water is 1 gm cm^{-3} and water offers negligible drag force on the bubble. The mass of the bubble is:

- (g = 980 cm/s^2).
 (A) 4.51 gm (B) 1.52 gm
 (C) 3.15 gm (D) 4.15 gm


17. Starting from the origin at time $t=0$, with initial velocity $5\hat{j} \text{ ms}^{-1}$, a particle moves in the $x - y$ plane with a constant acceleration of $(10\hat{i} + 4\hat{j}) \text{ ms}^{-2}$. At time t , its coordinates are $(20 \text{ m}, y_0 \text{ m})$. The values of t and y_0 are, respectively:

- (A) 2 s and 24 m (B) 4 s and 52 m
 (C) 5 s and 25 m (D) 2 s and 18 m

18. Two charged thin infinite plane sheets of uniform surface charge density σ_+ and σ_- , where $|\sigma_+| > |\sigma_-|$, intersect at right angle. Which of the following best represents the electric field lines for this system.




19. Blocks of masses m , $2m$ and $8m$ are arranged in a line on a frictionless floor. Another block of mass m , moving with speed v along the same line (see figure) collides with mass m in perfectly inelastic manner. All the subsequent collisions are also perfectly inelastic. By the time the last block of mass $8m$ starts moving the total energy loss is $p\%$ of the original energy. Value of 'p' is close to:




- (A) 77
(B) 37
(C) 87
(D) 94
20. A battery of 3.0 V is connected to a resistor dissipating 0.5 W of power. If the terminal voltage of the battery is 2.5 V , the power dissipated within the internal resistance is:
(A) 0.50 W
(B) 0.10 W
(C) 0.125 W
(D) 0.072 W
21. A closed vessel contains 0.1 mole of a monatomic ideal gas at 200 K . If 0.05 mole of the same gas at 400 K is added to it, the final equilibrium temperature (in K) of the gas in the vessel will be close to _____.
22. In a compound microscope, the magnified virtual image is formed at a distance of 25 cm from the eye-piece. The focal length of its objective lens is 1 cm . If the magnification is 100 and the tube length of the microscope is 20 cm , then the focal length of the eye-piece lens (in cm) is _____.

23. ABC is a plane lamina of the shape of an equilateral triangle. D, E are mid points of AB, AC and G is the centroid of the lamina. Moment of inertia of the lamina about an axis passing through G and perpendicular to the plane ABC is I_0 . If part ADE is removed, the moment of inertia of the remaining part about the same axis is $\frac{NI_0}{16}$ where N is an integer. Value of N is _____.




24. In the line spectra of hydrogen atom, difference between the largest and the shortest wavelengths of the Lyman series is $304\overset{\circ}{\text{Å}}$. The corresponding difference for the Paschan series in $\overset{\circ}{\text{Å}}$ is: _____.
25. A circular disc of mass M and radius R is rotating about its axis with angular speed ω_1 . If another stationary disc having radius $\frac{R}{2}$ and same mass M is dropped co-axially on to the rotating disc. Gradually both discs attain constant angular speed ω_2 . The energy lost in the process is p% of the initial energy. Value of p is _____.


PART – B (CHEMISTRY)

26. Among statements (a) – (d), the correct ones are:
 (a) Lime stone is decomposed to CaO during the extraction of iron from its oxides.
 (b) In the extraction of silver, silver is extracted as an anionic complex.
 (c) Nickel is purified by Mond's process.
 (d) Zr and Ti are purified by Van Arkel method.
 (A) (a), (b), (c) and (d) (B) (c) and (d) only
 (C) (a), (c) and (d) only (D) (b), (c) and (d) only
27. The decreasing order of reactivity of the following organic molecules towards AgNO₃ solution is:
- 


(A)




(B)




(C)




(D)
- (A) (A) > (B) > (D) > (C) (B) (C) > (D) > (A) > (B)
 (C) (A) > (B) > (C) > (D) (D) (B) > (A) > (C) > (D)
28. The region in the electromagnetic spectrum where the Balmer series lines appear is:
 (A) Ultraviolet (B) Visible
 (C) Infrared (D) Microwave
29. The pair in which both the species have the same magnetic moment (spin only) is:
 (A) [Co(OH)₄]²⁻ and [Fe(NH₃)₆]²⁺ (B) [Mn(H₂O)₆]²⁺ and [Cr(H₂O)]²⁺
 (C) [Cr(H₂O)₆]²⁺ and [Fe(H₂O)₆]²⁺ (D) [Cr(H₂O)₆]²⁺ and [CoCl₄]²⁻
30. On heating, lead (II) nitrate gives a brown gas (A). The gas (A) on cooling changes to a colourless solid/liquid (B). (B) on heating with NO changes to a blue solid (C). The oxidation number of nitrogen in solid (C) is:
 (A) +4 (B) +3
 (C) +2 (D) +5
31. [P] on treatment with Br₂/FeBr₃ in CCl₄ produced a single isomer C₈H₇O₂Br while heating [P] with sodalime gave toluene. The compound [P] is:
- 


(A)



(B)




(C)




(D)
32. The ionic radii of O²⁻, F⁻, Na⁺ and Mg²⁺ are in the order:
 (A) O²⁻ > F⁻ > Mg²⁺ > Na⁺ (B) Mg²⁺ > Na⁺ > F⁻ > O²⁻
 (C) F⁻ > O²⁻ > Na⁺ > Mg²⁺ (D) O²⁻ > F⁻ > Na⁺ > Mg²⁺


33. The intermolecular potential energy for the molecules A, B, C and D given below suggest that:



- (A) A – A has the largest bond enthalpy
 (B) A – D has the shortest bond length
 (C) D is more electronegative than other atoms
 (D) A – B has the stiffest bond.
34. An organic compound (A) (molecular formula $C_6H_{12}O_2$) was hydrolysed with dil. H_2SO_4 to give a carboxylic acid (B) and an alcohol (C). 'C' gives white turbidity immediately when treated with anhydrous $ZnCl_2$ and conc. HCl . The organic compound (A) is:




35. For the equilibrium $A \rightleftharpoons B$, the variation of the rate of the forward (a) and reverse (B) reaction with time is given by:




36. The number of isomers possible for $[\text{Pt}(\text{en})(\text{NO}_2)_2]$ is:
 (A) 4 (B) 3
 (C) 1 (D) 2

37. The IUPAC name of the following compound is:



- (A) 4-Bromo-5-methylcyclopentanoic acid
 (B) 4-Bromo-2-methylcyclopentane carboxylic acid
 (C) 3-Bromo-5-methylcyclopentane carboxylic acid
 (D) 5-Bromo-3-methylcyclopentanoic acid
38. What are the functional groups present in the structure of maltose?
 (A) One acetal and one hemiacetal (B) One ketal and one hemiketal
 (C) One acetal and one ketal (D) Two acetals

- 39.



$$E_{\text{Cu}^{2+}|\text{Cu}}^{\circ} = +0.34 \text{ V}$$

$$E_{\text{Zn}^{2+}|\text{Zn}}^{\circ} = -0.76 \text{ V}$$

Identify the incorrect statement from the options below for the above cell:

- (A) If $E_{\text{ext}} < 1.1 \text{ V}$, Zn dissolves at anode and Cu deposits at cathode
 (B) If $E_{\text{ext}} = 1.1 \text{ V}$, no flow of e^- or current occurs
 (C) If $E_{\text{ext}} > 1.1 \text{ V}$, Zn dissolves at Zn electrode and Cu deposits at Cu electrode
 (D) If $E_{\text{ext}} > 1.1 \text{ V}$, e^- flows from Cu to Zn
40. Which of the following will react with $\text{CHCl}_3 + \text{alc. KOH}$?
 (A) Adenine and proline (B) Adenine and lysine
 (C) Adenine and thymine (D) Thymine and proline

41. Match the following:

Column I		Column II	
(i)	Foam	(a)	smoke
(ii)	Gel	(b)	cell fluid
(iii)	Aerosol	(c)	jellies
(iv)	Emulsion	(d)	rubber
		(e)	froth
		(f)	milk

(A) (i)–(d), (ii)–(b), (iii)–(a), (iv)–(e)

(B) (i)–(d), (ii)–(b), (iii)–(e), (iv)–(f)

(C) (i)–(b), (ii)–(c), (iii)–(e), (iv)–(d)

(D) (i)–(e), (ii)–(c), (iii)–(a), (iv)–(f)

42. For one mole of an ideal gas, which of these statements must be true?

(a) U and H each depends only on temperature

(b) Compressibility factor z is not equal to 1

(c) $C_{P,m} - C_{V,m} = R$

(d) $dU = C_V dT$ for any process

(A) (a), (c) and (d)

(B) (b), (c) and (d)

(C) (a) and (c)

(D) (c) and (d)

43. On combustion of Li, Na and K in excess of air, the major oxides formed, respectively, are:





(A) Li_2O , Na_2O_2 and K_2O

(B) Li_2O , Na_2O_2 and KO_2

(C) Li_2O , Na_2O and K_2O_2

(D) Li_2O , Na_2O_2 and K_2O_2

44. When neopentyl alcohol is heated with an acid, it slowly converted into an 85:15 mixture of alkenes A and B, respectively. What are these alkenes?



45. The elements with atomic numbers 101 and 104 belong to, respectively:

(A) Actinoids and Group 6

(B) Actinoids and Group 4

(C) Group 11 and Group 4

(D) Group 6 and Actinoids



46. At 300K, the vapour pressure of a solution containing 1 mole of n-hexane and 3 moles of n-heptane is 550 mm of Hg. At the same temperature, if one more mole of n-heptane is added to this solution, the vapour pressure of the solution increases by 10 mm of Hg. What is the vapour pressure in mm Hg of n-heptane in its pure state _____?

47. If 75% of a first order reaction was completed in 90 minutes, 60% of the same reaction would be completed in approximately (in minutes) _____.

(Take: $\log 2 = 0.30$; $\log 2.5 = 0.40$)

48. The mass of ammonia in grams produced when 2.8 kg of dinitrogen quantitatively reacts with 1 kg of dihydrogen is _____.

49. The number of chiral centres present in [B] is _____.



50. At 20.0 mL solution containing 0.2 g impure H_2O_2 reacts completely with 0.316 g of KMnO_4 in acid solution. The purity of H_2O_2 (in %) is _____.
(mol. wt. of $\text{H}_2\text{O}_2 = 34$; mol. wt. of $\text{KMnO}_4 = 158$)

PART-C (MATHEMATICS)

51. Let α and β be roots of $x^2 - 3x + p = 0$ and γ and δ be the roots of $x^2 - 6x + q = 0$. If $\alpha, \beta, \gamma, \delta$, form a geometric progression. Then ratio $(2q + p) : (2q - p)$ is:
 (A) 33 : 31 (B) 5 : 3
 (C) 3 : 1 (D) 9 : 7
52. Let $f(x) = |x - 2|$ and $g(x) = f(f(x))$, $x \in [0, 4]$. Then $\int_0^3 (g(x) - f(x)) dx$ is equal to:
 (A) $\frac{1}{2}$ (B) $\frac{3}{2}$
 (C) 1 (D) 0
53. Let f be a twice differentiable function on $(1, 6)$. If $f(2) = 8$, $f'(2) = 5$, $f'(x) \geq 1$ and $f''(x) \geq 4$, for all $x \in (1, 6)$, then:
 (A) $f(5) + f'(5) \leq 26$ (B) $f(5) \leq 10$
 (C) $f(5) + f'(5) \geq 28$ (D) $f'(5) + f''(5) \leq 20$
54. Let $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ ($a > b$) be given ellipse, length of whose latus rectum is 10. If its eccentricity is the maximum value of the function, $\phi(t) = \frac{5}{12} + t - t^2$, then $a^2 + b^2$ is equal to:
 (A) 126 (B) 135
 (C) 116 (D) 145
55. The mean and variance of 8 observations are 10 and 13.5, respectively. If 6 of these observations are 5, 7, 10, 12, 14, 15, then the absolute difference of the remaining two observations is:
 (A) 3 (B) 5
 (C) 7 (D) 9
56. Let x_0 be the point of local maxima of $f(x) = \vec{a} \cdot (\vec{b} \times \vec{c})$ where $\vec{a} = x\hat{i} - 2\hat{j} + 3\hat{k}$, $\vec{b} = -2\hat{i} + x\hat{j} - \hat{k}$ and $\vec{c} = 7\hat{i} - 2\hat{j} + x\hat{k}$. Then the value of $\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}$ at $x = x_0$ is
 (A) -4 (B) -22
 (C) -30 (D) 14
57. If $(a + \sqrt{2}b \cos x)(a - \sqrt{2}b \cos y) = a^2 - b^2$, where $a > b > 0$, then $\frac{dx}{dy}$ at $\left(\frac{\pi}{4}, \frac{\pi}{4}\right)$ is:
 (A) $\frac{a - 2b}{a + 2b}$ (B) $\frac{2a + b}{2a - 2}$
 (C) $\frac{a + b}{a - b}$ (D) $\frac{a - b}{a + b}$
58. Two vertical poles AB = 15 m and CD = 10 m are standing apart on a horizontal ground with points A and C on the ground. If P is the point of intersection of BC and AD, then the height of P (in m) above the line AC is:
 (A) $\frac{10}{3}$ (B) 6
 (C) 5 (D) $\frac{20}{3}$

59. If $1 + (1 - 2^2 \cdot 1) + (1 - 4^2 \cdot 3) + (1 - 6^2 \cdot 5) + \dots + (1 - 20^2 \cdot 19) = \alpha - 220\beta$, then an ordered pair (α, β) is equal to:
 (A) (11, 97) (B) (10, 97)
 (C) (10, 103) (D) (11, 103)
60. Given the following two statements:
 $(S_1): (q \vee p) \rightarrow (p \leftrightarrow \sim q)$ is a tautology.
 $(S_2): \sim q \wedge (\sim p \leftrightarrow q)$ is a fallacy. Then:
 (A) both (S_1) and (S_2) are correct (B) only (S_2) is correct
 (C) both (S_1) and (S_2) are incorrect (D) only (S_1) is correct
61. Let $u = \frac{2z+i}{z-ki}$, $z = x+iy$ and $k > 0$. If the curve represented by $\text{Re}(u) + \text{Im}(u) = 1$ intersects the y-axis at the points P and Q where $PQ = 5$, then the value of k is:
 (A) 2 (B) $\frac{1}{2}$
 (C) $\frac{3}{2}$ (D) 4
62. A triangle ABC lying in the first quadrant has two vertices as A(1, 2) and B(3, 1). If $\angle BAC = 90^\circ$, and $\text{ar}(\triangle ABC) = 5\sqrt{5}$ sq. units, then the abscissa of the vertex C is:
 (A) $1+2\sqrt{5}$ (B) $1+\sqrt{5}$
 (C) $2\sqrt{5}-1$ (D) $2+\sqrt{5}$
63. Let P (3, 3) be a point on the hyperbola, $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$. If the normal to it at P intersects the x-axis at (9, 0) and e is its eccentricity, then the ordered pair (a^2, e^2) is equal to:
 (A) $\left(\frac{3}{2}, 2\right)$ (B) $\left(\frac{9}{2}, 2\right)$
 (C) (9, 3) (D) $\left(\frac{9}{2}, 3\right)$
64. The integral $\int \left(\frac{x}{x \sin x + \cos x}\right)^2 dx$ is equal to (where C is constant of integration):
 (A) $\tan x + \frac{x \sec x}{x \sin x + \cos x} + C$ (B) $\tan x - \frac{x \sec x}{x \sin x + \cos x} + C$
 (C) $\sec x + \frac{x \tan x}{x \sin x + \cos x} + C$ (D) $\sec x - \frac{x \tan x}{x \sin x + \cos x} + C$
65. If $A = \begin{bmatrix} \cos \theta & i \sin \theta \\ i \sin \theta & \cos \theta \end{bmatrix}$, $\left(\theta = \frac{\pi}{24}\right)$ and $A^5 = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, where $i = \sqrt{-1}$, then, which one of the following is not true?
 (A) $a^2 - c^2 = 1$ (B) $a^2 - b^2 = \frac{1}{2}$
 (C) $0 \leq a^2 + b^2 \leq 1$ (D) $a^2 - d^2 = 0$

66. The value of $\sum_{r=0}^{20} {}^{50-r}C_6$ is equal to:
 (A) ${}^{50}C_6 - {}^{30}C_6$ (B) ${}^{51}C_7 + {}^{30}C_7$
 (C) ${}^{51}C_7 - {}^{30}C_7$ (D) ${}^{50}C_7 - {}^{30}C_7$
67. Let $[t]$ denote the greatest integer $\leq t$. Then the equation in x , $[x]^2 + 2[x + 2] - 7 = 0$ has:
 (A) infinitely many solutions (B) exactly two solutions
 (C) no integral solution (D) exactly four integral solutions
68. Let $f(x) = \int \frac{\sqrt{x}}{(1+x)^2} dx$ ($x \geq 0$). Then $f(3) - f(1)$ is equal to:
 (A) $\frac{\pi}{12} + \frac{1}{2} - \frac{\sqrt{3}}{4}$ (B) $-\frac{\pi}{12} + \frac{1}{2} + \frac{\sqrt{3}}{4}$
 (C) $\frac{\pi}{6} + \frac{1}{2} - \frac{\sqrt{3}}{4}$ (D) $-\frac{\pi}{6} + \frac{1}{2} + \frac{\sqrt{3}}{4}$
69. A survey shows that 63% of the people in a city read newspaper A whereas 76% read newspaper B. if $x\%$ of the people read both the newspapers, then a possible value of x can be:
 (A) 29 (B) 55
 (C) 37 (D) 65
70. Let $y = y(x)$ be the solution of the differential equation, $xy' - y = x^2(x \cos x + \sin x)$, $x > 0$. If $y(\pi) = \pi$, then $y''\left(\frac{\pi}{2}\right) + y\left(\frac{\pi}{2}\right)$ is equal to:
 (A) $1 + \frac{\pi}{2} + \frac{\pi^2}{4}$ (B) $2 + \frac{\pi}{2} + \frac{\pi^2}{4}$
 (C) $2 + \frac{\pi}{2}$ (D) $1 + \frac{\pi}{2}$
71. Let $(2x^2 + 3x + 4)^{10} = \sum_{r=0}^{20} a_r x^r$. Then $\frac{a_7}{a_{13}}$ is equal to _____.
72. If the equation of a plane P, passing through the intersection of the planes, $x + 4y - z + 7 = 0$ and $3x + y + 5z = 8$ is $ax + by + 6z = 15$ for some $a, b \in \mathbb{R}$, then the distance of the point $(3, 2, -1)$ from the plane P is _____.
73. If the system of equations
 $x - 2y + 3z = 9$
 $2x + y + z = b$
 $x - 7y + az = 24$,
 has infinitely many solutions, then $a - b$ is equal to _____.
74. Suppose a differentiable function $f(x)$ satisfies the identity $f(x + y) = f(x) + f(y) + xy^2 + x^2y$, for all real x and y . If $\lim_{x \rightarrow 0} \frac{f(x)}{x} = 1$ then $f'(3)$ is equal to _____.
75. The probability of a man hitting a target is $\frac{1}{10}$. The least number of shots required, so that the probability of his hitting the target at least once is greater than $\frac{1}{4}$, is _____.